

A NONCOMMUTATIVE VERSION OF THE JOHN-NIRENBERG THEOREM

MARIUS JUNGE⁽¹⁾ AND MAGDALENA MUSAT⁽²⁾

ABSTRACT. We prove a noncommutative version of the John-Nirenberg theorem for non-tracial filtrations of von Neumann algebras. As an application, we obtain an analogue of the classical large deviation inequality for elements of the associated BMO space.

1. INTRODUCTION

The John-Nirenberg theorem is an important tool in analysis and probability. It provides a characterization of BMO , the space of functions of *bounded mean oscillation*. In its first version it was proved by John and Nirenberg [13] in 1961. Through its connection with the theory of Hardy spaces, the John-Nirenberg result has many applications in harmonic analysis and the theory of singular integrals (see, e.g., Stein [37, 38]). We refer to the books of Bennett and Sharpley [5], Garcia-Cuerva and Rubio de Francia [9], Garnett [10] and Koosis [22] for the interval (function space) version, respectively to the work of Bass [1], Garsia [11] and Petersen [29] for the martingale version of this theorem.

In this paper we analyze analogues of the John-Nirenberg results in a noncommutative setting. We first recall the classical results. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $(\mathcal{F}_n)_{n \geq 0}$ an increasing sequence of sub- σ -algebras of $\mathcal{F}$. For $1 \leq p < \infty$ consider the norms

$$(1.1) \quad \|x\|_{BMO_p} := \sup_n \|\mathbb{E}(|x - x_{n-1}|^p | \mathcal{F}_n)\|_{\infty}^{\frac{1}{p}},$$

where $x_n = \mathbb{E}(x | \mathcal{F}_n)$, for all non-negative integers n . The usual BMO norm corresponds to $p = 2$ above, i.e., $\|x\|_{BMO} = \|x\|_{BMO_2}$. The John-Nirenberg theorem yields universal constants $C_1, C_2 > 0$ with the following property. If $\|x\|_{BMO_1} < C_2$, then

$$(1.2) \quad \sup_n \|\mathbb{E}(e^{C_1|x-x_{n-1}|} | \mathcal{F}_n)\|_{\infty} < 1.$$

Using the power series expansion for the exponential function, it follows that (1.2) is equivalent to

$$(1.3) \quad \|\mathbb{E}(|x - x_{n-1}|^p | \mathcal{F}_n)\|_{\infty}^{\frac{1}{p}} \leq Cp \|x\|_{BMO},$$

[†]Junge is partially supported by the National Science Foundation, DMS-0301116.

2000 *Mathematics Subject Classification.* Primary: 46L52; Secondary: 60G46.

Key words and phrases. noncommutative L_p -spaces and BMO ; noncommutative martingales; interpolation.

for all $n \geq 0$ and all $1 \leq p < \infty$, where $C > 0$ is a universal constant. Furthermore, a standard duality argument yields the equality

$$(1.4) \quad \|\mathbb{E}(|x - x_{n-1}|^p | \mathcal{F}_n)\|_\infty^{\frac{1}{p}} = \sup_{a \in L_p(\mathcal{F}_n), \|a\|_p \leq 1} \|(x - x_{n-1})a\|_p.$$

It can be easily seen that condition (1.3) implies that, for $1 \leq p < \infty$,

$$(1.5) \quad \|x\|_p \leq Cp \|x\|_{BMO}.$$

Moreover, the John-Nirenberg theorem follows from the validity of (1.5) for arbitrary probability measures and filtrations. Indeed, let $1 \leq p < \infty$, and fix an integer $n \geq 0$. Consider a positive element $a \in L_p(\mathcal{F}_n)$ with $\|a\|_p = 1$, and denote by μ_a the probability measure defined by $d\mu_a = a^{\frac{1}{p}} d\mathbb{P}$. Then

$$(1.6) \quad \|(x - x_{n-1})a\|_p = \|x - x_{n-1}\|_{L_p(\mu_a)}.$$

Denote by $BMO((\mathcal{F}_k)_{k \geq n}, \mu_a)$ the BMO space associated to the triplet $(\Omega, \mathcal{F}, \mu_a)$ and the filtration $(\mathcal{F}_k)_{k \geq n}$ of $\mathcal{F}$. Applying (1.5) for the probability measure μ_a and the new filtration $(\mathcal{F}_k)_{k \geq n}$ of $\mathcal{F}$, it follows that

$$(1.7) \quad \|x - x_{n-1}\|_{L_p(\mu_a)} \leq Cp \|x - x_{n-1}\|_{BMO((\mathcal{F}_k)_{k \geq n}, \mu_a)}.$$

Note that $\|x - x_{n-1}\|_{BMO((\mathcal{F}_k)_{k \geq n}, \mu_a)} \leq \|x\|_{BMO((\mathcal{F}_k)_{k \geq 0}, \mathbb{P})}$. An application of (1.6), together with formula (1.4) yields now the inequality (1.3).

In the form (1.5), one can formulate a noncommutative version of the John-Nirenberg theorem. First this requires an appropriate definition of BMO spaces. In their seminal paper [32], Pisier and Xu proved the noncommutative analogues of the Burkholder-Gundy square function inequalities. Fermionic versions of the square function inequalities had previously been considered by Carlen and Krée [4]. In [32], Pisier and Xu also introduced the BMO space for noncommutative martingales and proved the analogue of the classical Fefferman-Stein duality $BMO = (H_1)^*$. Their work triggered a rapid development in the L_p -theory of noncommutative martingales, see, e.g., Junge [14], Junge and Xu [18, 19, 20] and Randrianantoanina [34, 35].

In the following we assume that $\mathcal{N}$ is a von Neumann algebra, ϕ a normal faithful state on $\mathcal{N}$ and $(\mathcal{N}_n)_{n \geq 0}$ an increasing sequence of von Neumann subalgebras whose union generates $\mathcal{N}$ in the w^* -topology. Moreover, we assume that for all positive integers n there exist normal conditional expectations $\mathcal{E}_n : \mathcal{N} \rightarrow \mathcal{N}_n$ such that

$$(1.8) \quad \phi(\mathcal{E}_n(x)y) = \phi(xy)$$

for all $x \in \mathcal{N}$ and $y \in \mathcal{N}_n$. We recall the following definitions (see [18], [32])

$$\|x\|_{BMO^c} = \sup_m \sup_{n \leq m} \|\mathcal{E}_n((x_m - x_{n-1})^*(x_m - x_{n-1}))\|_\infty^{\frac{1}{2}},$$

where $x_n = \mathcal{E}_n(x)$, for all $n \geq 0$. Respectively,

$$\|x\|_{BMO} = \max\{\|x\|_{BMO^c}, \|x^*\|_{BMO^c}\}.$$

The conditioned L_∞ -spaces, $L_\infty^c(\mathcal{N}, \mathcal{E}_n)$ associated to the conditional expectations $\mathcal{E}_n$ were introduced in [14] (see also [36]) by defining

$$\|x\|_{L_\infty^c(\mathcal{N}, \mathcal{E}_n)} = \|\mathcal{E}_n(x^*x)\|_\infty^{\frac{1}{2}}.$$

We shall point out that in general, for $2 < p < \infty$, the expression $\|\mathcal{E}_n((x^*x)^{\frac{p}{2}})\|_\infty^{\frac{1}{p}}$ does not necessarily provide a norm in the noncommutative setting. Therefore, we will use a different approach to generalize the BMO_p -norms. Namely, motivated by (1.4), we define for $2 \leq p \leq \infty$,

$$(1.9) \quad \|x\|_{BMO_p^c} = \sup_m \sup_{n \leq m} \sup_{a \in L_p(N_n), \|a\|_p \leq 1} \|(x_m - x_{n-1})a\|_p.$$

Respectively, we define

$$(1.10) \quad \|x\|_{BMO_p} = \max\{\|x\|_{BMO_p^c}, \|x^*\|_{BMO_p^c}\}.$$

These norms can, in fact, be obtained by interpolation between conditional L_p -spaces. More details are given in Section 4. (See [16] for a more general discussion of such norms.)

Our martingale version of the John-Nirenberg theorem reads as follows.

Theorem 1.1. *There exists a universal constant $c > 0$ such that for all $2 < p < \infty$,*

$$(1.11) \quad \|x\|_{BMO} \leq \|x\|_{BMO_p} \leq cp \|x\|_{BMO}.$$

The order cp of the constant in the inequality (1.11) above is the same as in the commutative setting. In a preliminary version of this paper we were only able to prove that (1.11) holds with a constant of the order cp^2 . The right order cp was obtained using very recent results of Randrianantoanina [35] on the optimal order of growth for the constants in the noncommutative Burkholder-Gundy square function inequalities. For the proof of Theorem 1.1, note that formula (1.9), which is the key point in the change of density argument described in classical setting, leads to the passage from traces to various states on the von Neumann algebra. Therefore we consider martingales with respect to states and their modular theory.

As an application of a noncommutative version of Chebychev inequality proved by Defant and Junge [8], we obtain an analogue of the classical large deviation inequality for elements of BMO . Namely, if $\|x\|_{BMO} < 1$, then

$$(1.12) \quad \mathbb{P}\{|x - x_0| > t\} < C_2 e^{-tC_1}.$$

for all $t > 0$, where C_1, C_2 are universal constants (see [10], [11]). More precisely, we prove

Theorem 1.2. *There exist universal constants $c_1, c_2 > 0$ such that if $\|x\|_{BMO} < 1$, then for all $t > 0$, there exists a projection $f \in \mathcal{N}$ such that $\|(x - x_0)f\| \leq t$ and*

$$(1.13) \quad \phi(1 - f) < c_2 e^{-tc_1}.$$

In the commutative case $f = 1_A$, for some measurable subset A of Ω . The condition $\|(x - x_0)f\| \leq t$ implies that $A \subseteq \{\omega \in \Omega : |x(\omega) - x_0(\omega)| \leq t\}$. Then (1.13) yields (1.12).

Our paper is organized as follows. In section 2, we explain the notation and Kosaki's interpolation results, a fundamental tool in our argument. In section 3, we discuss various inclusions of BMO into L_p . Note that even the inclusion $BMO \subset L_2$ is not immediate in the non-tracial setting. We first establish that for $2 \leq p < \infty$, (BMO, L_p) forms an interpolation couple and prove that $[BMO, L_p]_{\frac{p}{p-2}} = L_q$, $2 \leq p < q < \infty$ (extending the results in [26] to the non-tracial setting). By general theory of interpolation, the continuous inclusion $L_q \subset L_p$ yields a continuous inclusion $BMO \subset L_p$. The inequality (1.11) and the modifications required to deduce the interval version of the John-Nirenberg theorem are discussed in section 4, as well as the proof of Theorem 1.2. We are indebted to Tao Mei and Narcisse Randrianantoanina for providing us with the preprints [25], respectively [35].

2. PRELIMINARIES

We use standard notation in operator algebras. We refer to [21] and [41] for background on von Neumann algebras. In the following, we will consider a σ -finite von Neumann algebra $\mathcal{N}$ acting on a Hilbert space H , and a distinguished normal faithful state ϕ on $\mathcal{N}$. We denote by $\sigma_t = \sigma_t^\phi$ the one parameter modular automorphism group on $\mathcal{N}$ associated with ϕ . Haagerup's abstract L_p -spaces are defined using the crossed-product $\mathcal{R} = \mathcal{N} \rtimes_{\sigma_t} \mathbb{R}$. We recall that $\mathcal{R} \subseteq \mathcal{B}(L_2(\mathbb{R}, H))$ is the von Neumann algebra generated by the operators $\pi(x)$, $x \in \mathcal{N}$ and, respectively, $\lambda(s)$, $s \in \mathbb{R}$, where

$$\pi(x)(\xi)(t) = \sigma_{-t}(x)\xi(t) \quad \text{and} \quad \lambda(t)(\xi)(s) = \xi(t-s),$$

for all $\xi \in \mathcal{B}(L_2(\mathbb{R}, H))$ and all $t \in \mathbb{R}$. In the following, we may and will identify $\mathcal{N}$ with $\pi(\mathcal{N})$, since π is a normal faithful representation of $\mathcal{N}$ on $L_2(\mathbb{R}, H)$. Furthermore, note that π is invariant under the dual action

$$\theta_s(x) = W(s)xW(s)^*, \quad s \in \mathbb{R}, x \in \mathcal{R},$$

and, moreover,

$$\pi(\mathcal{N}) = \{x \in \mathcal{R} : \theta_s(x) = x, \text{ for all } s \in \mathbb{R}\}.$$

Here the unitary operators $W(s)$, $s \in \mathbb{R}$ are defined by the phase shift

$$W(s)(\xi)(t) = e^{-ist}\xi(t), \quad t \in \mathbb{R}.$$

As shown in [28], the crossed product $\mathcal{N} \rtimes_{\sigma_t} \mathbb{R}$ is semifinite and admits a unique normal semifinite trace τ such that for all $s \in \mathbb{R}$,

$$\tau(\theta_s(x)) = e^{-s}\tau(x).$$

For $1 \leq p \leq \infty$, $L_p(\mathcal{N})$ is defined as the space of all τ -measurable operators x affiliated with $\mathcal{R} = \mathcal{N} \rtimes_{\sigma_t} \mathbb{R}$ such that for all $s \in \mathbb{R}$,

$$\theta_s(x) = e^{-s/p}x.$$

It follows from the definition that $L_\infty(\mathcal{N})$ coincides with $\mathcal{N}$. Furthermore, there is a canonical isomorphism between $L_1(\mathcal{N})$ and the predual $\mathcal{N}_*$ of $\mathcal{N}$. This requires some explanation. Following [28], every normal semifinite faithful weight (n.s.f., for short) $\psi \in (\mathcal{N}_*)_+$ is given by a density $h_\psi \in L_1(\mathcal{N})_+$ satisfying

$$\tau(h_\psi x) = \int_{\mathbb{R}} \psi(\theta_s(x)) \, ds,$$

for all $x \in \mathcal{R}_+$. Using the polar decomposition of an arbitrary element $\psi \in \mathcal{N}_*$, this correspondence between $(\mathcal{N}_*)_+$ and $L_1(\mathcal{N})_+$ extends to a bijection between $\mathcal{N}_*$ and $L_1(\mathcal{N})$. Namely, if $\psi \in \mathcal{N}_*$, then $\psi = u|\psi|$, where $u \in \mathcal{N}$ and $|\psi|$ is the modulus of ψ . By construction, the corresponding $h_\psi \in L_1(\mathcal{N})$ admits the polar decomposition

$$h_\psi = u|h_\psi| = uh_{|\psi|}.$$

We may define a norm on $L_1(\mathcal{N})$ by

$$\|h_\psi\|_1 = |\psi|(1) = \|\psi\|_{\mathcal{N}_*}, \quad \psi \in \mathcal{N}_*.$$

In this way we obtain the isometry between $L_1(\mathcal{N})$ and $\mathcal{N}_*$. Furthermore, define a linear functional $\text{tr} : L_1(\mathcal{N}) \rightarrow \mathbb{C}$, called *trace*, by

$$\text{tr}(h_\psi) = \psi(1).$$

It is important to note that if $\frac{1}{p} + \frac{1}{q} = 1$, then for all $x \in L_p(\mathcal{N})$ and $y \in L_q(\mathcal{N})$, we have the tracial property

$$\text{tr}(xy) = \text{tr}(yx).$$

Let us now return to the special state ϕ fixed at the beginning. Then, as explained above, there exists a density $D_\phi \in L_1(\mathcal{N})$ such that for all $x \in \mathcal{N}$,

$$\phi(x) = \text{tr}(D_\phi x)$$

In the following we drop the subscript and reserve the letter D exclusively for this density. Given $1 \leq p < \infty$ and $x \in L_p(\mathcal{N})$, define

$$\|x\|_p = (\text{tr}(|x|^p))^{1/p}, \quad \text{respectively} \quad \|x\|_\infty = \|x\|_{\mathcal{N}}.$$

If $\frac{1}{p} + \frac{1}{q} = \frac{1}{r}$, then for all $x \in L_p(\mathcal{N})$ and $y \in L_q(\mathcal{N})$, Hölder's inequality holds, i.e.,

$$\|xy\|_r \leq \|x\|_p \|y\|_q.$$

As a consequence, given $1 \leq p < \infty$, the mapping $(x, y) \in L_p(\mathcal{N}) \times L_{p'}(\mathcal{N}) \mapsto \text{tr}(xy)$, where $\frac{1}{p} + \frac{1}{p'} = 1$, defines a duality bracket between $L_p(\mathcal{N})$ and $L_{p'}(\mathcal{N})$, with respect to which we have the isometry

$$(L_p(\mathcal{N}))^* = L_{p'}(\mathcal{N}).$$

In the sequel we will make repeated use of the following well-known fact.

Fact 2.1. *Let $1 \leq p, q, r \leq \infty$ such that $\frac{1}{p} = \frac{1}{q} + \frac{1}{r}$ and $p < \infty$. Let $x \in L_q(\mathcal{N})$, and $a \in L_r(\mathcal{N})$, $b \in L_r(\mathcal{N})$ be positive elements. Then the map*

$$f(z) = a^{1-z}xb^z \in L_p(\mathcal{N})$$

is continuous on $S = \{z \in \mathbb{C} : 0 \leq \operatorname{Re}(z) \leq 1\}$ and analytic in the interior S_0 of S .

Consider now the 1-parameter automorphism group

$$\alpha_t(x) = D^{it}xD^{-it}, \quad t \in \mathbb{R}.$$

Note that α_t is strongly continuous and leaves $\mathcal{N}$ invariant since

$$\theta_s(D^{it}xD^{-it}) = e^{its}D^{it}xe^{-its}D^{its} = D^{it}xD^{-it}.$$

From Fact 2.1 we deduce that for every $x \in \mathcal{N}$, the map $f(z) = D^{1-z}xD^z$ is analytic, taking values in $L_1(\mathcal{N})$. In particular, it follows that the map $f_{x,y}(z) = \operatorname{tr}(D^{1-z}xD^zy)$ is analytic and satisfies

$$f_{x,y}(it) = \phi(x\alpha_t(y)) \quad \text{and} \quad f_{x,y}(1+it) = \phi(\alpha_t(y)x),$$

for all $t \in \mathbb{R}$. Since

$$\phi(\alpha_t(x)) = \operatorname{tr}(DD^{it}xD^{-it}) = \phi(x),$$

we deduce that $\sigma_t = \alpha_t$, for all $t \in \mathbb{R}$ (see, e.g., [39, Section 2.12]).

An element $x \in \mathcal{N}$ is called analytic if the map $t \in \mathbb{R} \mapsto \sigma_t(x) \in \mathcal{N}$ extends to an analytic function $z \in \mathbb{C} \mapsto \sigma_z(x) \in \mathcal{N}$. The family $\mathcal{N}_a$ of analytic elements in $\mathcal{N}$ is a weak*-dense *-subalgebra of $\mathcal{N}$ (see [28]). Given $1 \leq p < \infty$ and $0 \leq \theta \leq 1$, it can be shown (see, e.g., [14]) that the space $\mathcal{N}_a D^{1/p}$ is dense in $L_p(\mathcal{N})$ and

$$D^{\frac{1-\theta}{p}} \mathcal{N}_a D^{\frac{\theta}{p}} = \mathcal{N}_a D^{\frac{1}{p}}.$$

We will denote by $L_p(\mathcal{N})_a$ the family of elements with the property that the map $t \mapsto \sigma_t(x)$, $t \in \mathbb{R}$ extends to an analytic function on $\mathbb{C}$ with values in $L_p(\mathcal{N})$. Since $\mathcal{N}_a D^{\frac{1}{p}} \subset L_p(\mathcal{N})_a$ is dense in $L_p(\mathcal{N})$, as mentioned above, it follows that $L_p(\mathcal{N})_a$ is dense in $L_p(\mathcal{N})$.

Since interpolation is our main tool in the paper, we briefly recall in the following some basic notions concerning the complex method of interpolation due to Calderón. Our main reference for interpolation theory is Bergh and Löfström [3]. A pair of Banach spaces (X_0, X_1) is called a *compatible couple* if they embed continuously in some topological vector space X . This allows us to consider the spaces $X_0 \cap X_1$ and $X_0 + X_1$ (by identifying them inside X). They are Banach spaces when equipped, respectively, with the following norms

$$(2.14) \quad \|x\|_{X_0 \cap X_1} = \max\{\|x\|_{X_0}, \|x\|_{X_1}\},$$

$$(2.15) \quad \|x\|_{X_0 + X_1} = \inf\{\|x_0\|_{X_0} + \|x_1\|_{X_1} : x = x_0 + x_1, x_0 \in X_0, x_1 \in X_1\}.$$

Following the notation from [3], let $\mathcal{F}$ be the family of all continuous and bounded functions $f : S \rightarrow X_0 + X_1$ satisfying the following properties

- (1) f is analytic in S_0 ,
- (2) $f(it) \in X_0$ and $f(1+it) \in X_1$ for all $t \in \mathbb{R}$,
- (3) $f(it) \rightarrow 0$ and $f(1+it) \rightarrow 0$ as $t \rightarrow \infty$.

By the Phragmen-Lindelöf theorem, $\mathcal{F}$ is a Banach space under the norm

$$\|f\|_{\mathcal{F}} = \max \left\{ \sup_{t \in \mathbb{R}} \|f(it)\|_{X_0}, \sup_{t \in \mathbb{R}} \|f(1+it)\|_{X_1} \right\}.$$

For $0 \leq \theta \leq 1$, set $[X_0, X_1]_{\theta} = \{x \in X_0 + X_1 : x = f(\theta), \text{ for some } f \in \mathcal{F}\}$. This is called the *complex interpolation space* (of exponent θ) between X_0 and X_1 , and it is a Banach space under the norm

$$\|x\|_{\theta} = \inf \{\|f\|_{\mathcal{F}} : x = f(\theta), f \in \mathcal{F}\}.$$

The complex method is an exact interpolation functor; i.e., if $T : X_0 + X_1 \rightarrow Y_0 + Y_1$ is a linear operator which is bounded both from X_0 to Y_0 (with norm M_0) and from X_1 to Y_1 (with norm M_1), then T is bounded from $[X_0, X_1]_{\theta}$ to $[Y_0, Y_1]_{\theta}$, with norm $\leq M_0^{1-\theta} M_1^{\theta}$.

3. INTERPOLATION RESULTS

In this paper we will make crucial use of Kosaki's interpolation results (see [23]), which will enable us to extend the results in [26] to the non-tracial setting. For $1 \leq p < q \leq \infty$ and $0 \leq \eta \leq 1$ consider the mapping $I_{q,p}^{\eta} : L_q(\mathcal{N}) \rightarrow L_p(\mathcal{N})$ defined by

$$I_{q,p}^{\eta}(x) = D^{(1-\eta)(\frac{1}{p}-\frac{1}{q})} x D^{\eta(\frac{1}{p}-\frac{1}{q})}.$$

Note that $I_{q,p}^{\eta}$ is injective. In [23], Kosaki introduces the spaces

$$(3.16) \quad L_p^{\eta}(\mathcal{N}, \phi) = [I_{\infty,1}^{\eta}(\mathcal{N}), L_1(\mathcal{N})]_{\frac{1}{p}},$$

for $1 \leq p \leq \infty$. We should point out that Kosaki's original notation is slightly different, namely $L_p^{\eta}(\mathcal{N}, \phi)$ is denoted in [23] by $i_p^{\eta}(L_p)$, where i_p^{η} denotes the mapping $I_{p,1}^{\eta}$. Also, in our terminology the space $L_p^{\eta}(\mathcal{N}, \phi)$ is considered as a subspace of $L_1(\mathcal{N})$ whereas Kosaki formally works in $\mathcal{N}_{*}$. Using the canonical isometric isomorphism between $L_1(\mathcal{N})$ and $\mathcal{N}_{*}$ explained in the preliminaries, Kosaki's results may be reformulated in this language as

$$(3.17) \quad L_p^{\eta}(\mathcal{N}, \phi) = D^{\frac{1-\eta}{p'}} L_p(\mathcal{N}) D^{\frac{\eta}{p'}},$$

where $\frac{1}{p} + \frac{1}{p'} = 1$. This is not only an equality on the level of sets, but it means that if $x \in L_p(\mathcal{N})$ then $D^{\frac{1-\eta}{p'}} x D^{\frac{\eta}{p'}} \in [I_{\infty,1}^{\eta}(\mathcal{N}), L_1(\mathcal{N})]_{\frac{1}{p}}$ and, furthermore,

$$\|x\|_p = \left\| D^{\frac{1-\eta}{p'}} x D^{\frac{\eta}{p'}} \right\|_{[I_{\infty,1}^{\eta}(\mathcal{N}), L_1(\mathcal{N})]_{\frac{1}{p}}}.$$

Moreover, every element in the interpolation space $[I_{\infty,1}^\eta(\mathcal{N}), L_1(\mathcal{N})]_{\frac{1}{p}}$ comes from an element $x \in L_p(\mathcal{N})$. In our context, we will also consider the interpolation couple (A_0, A_1) , where $A_0 = I_{q,p}^\eta(L_q(\mathcal{N}))$ and $A_1 = L_p(\mathcal{N})$, with $1 \leq p < q \leq \infty$. We will use the notation

$$\|I_{q,p}^\eta(x)\|_{A_0} = \|x\|_q.$$

This is to emphasize that the ambient topological vector space for the interpolation couple $(I_{q,p}^\eta(L_q(\mathcal{N})), L_p(\mathcal{N}))$ is $L_p(\mathcal{N})$, and the inclusion map of $L_q(\mathcal{N})$ into $L_p(\mathcal{N})$ is given by $I_{q,p}^\eta$. Using (3.16), (3.17) and the reiteration theorem for complex interpolation ([3, Theorem 4.6.1]), we deduce for $1 \leq p < s < q \leq \infty$ and $0 < \theta < 1$ with $\frac{1}{s} = \frac{1-\theta}{q} + \frac{\theta}{p}$ that

$$(3.18) \quad I_{s,p}^\eta(L_s(\mathcal{N})) = [I_{q,p}^\eta(L_q(\mathcal{N})), L_p(\mathcal{N})]_\theta.$$

We will now use these interpolation spaces in the context of BMO-type norms. Indeed, it is helpful to consider the following vector-valued ℓ_∞ space defined for sequences $(x_n)_{n \geq 0}$ in $L_p(\mathcal{N})$ as follows

$$\|(x_n)\|_{L_p^c(\mathcal{N}; \ell_\infty)} = \left\| \sup_n (x_n^* x_n) \right\|_{\frac{p}{2}}^{\frac{1}{2}},$$

where the right hand side is to be understood in the sense of the following suggestive notation introduced in [14]. Namely, if $1 \leq q, q' \leq \infty$ with $\frac{1}{q} + \frac{1}{q'} = 1$ then, for a finite sequence of positive elements in $L_p(\mathcal{N})$, set

$$(3.19) \quad \left\| \sup_n x_n \right\|_q = \sup \left\{ \sum_{n \geq 0} \text{tr}(x_n y_n) : y_n \geq 0, \left\| \sum_{n \geq 0} y_n \right\|_{q'} \leq 1 \right\}.$$

In the commutative case this coincides with the usual definition of the norm in the l_∞ -valued L_q -space. Here the actual supremum does not necessarily make sense as in the commutative case and thus it is just a notation. Note that $L_p^c(\mathcal{N}, \ell_\infty)$ is only well-defined for $p \geq 2$.

Remark 3.1. Let $1 \leq p < q, s < \infty$ be such that $\frac{1}{p} = \frac{1}{q} + \frac{1}{s}$. Then, for all $0 \leq a, b \in L_s(\mathcal{N})$,

- 1) $\sup_m \left\| \sup_{n \leq m} x_n \right\|_p = \left\| \sup_n x_n \right\|_p$
- 2) $\left\| \sup_n a^{\frac{1}{2}} x_n a^{\frac{1}{2}} \right\|_p \leq \|a\|_s \left\| \sup_n x_n \right\|_q$
- 3) $\left\| \sup_n x_n^{\frac{1}{2}} b x_n^{\frac{1}{2}} \right\|_p \leq \|b\|_\infty \left\| \sup_n x_n \right\|_p.$

Proof. Item 1) is an immediate consequence of the definition (3.19). Item 2) follows from Hölder's inequality, namely,

$$\begin{aligned}
\left\| \sup_n a^{\frac{1}{2}} x_n a^{\frac{1}{2}} \right\|_p &= \sup \left\{ \sum_{n \geq 0} \operatorname{tr}(a^{\frac{1}{2}} x_n a^{\frac{1}{2}} y_n) : y_n \geq 0, \left\| \sum_{n \geq 0} y_n \right\|_{p'} \leq 1 \right\} \\
&= \sup \left\{ \sum_{n \geq 0} \operatorname{tr}(x_n a^{\frac{1}{2}} y_n a^{\frac{1}{2}}) : y_n \geq 0, \left\| \sum_{n \geq 0} y_n \right\|_{p'} \leq 1 \right\} \\
&\leq \|a\|_s \sup_n \|x_n\|_q.
\end{aligned}$$

Item 3) follows from definition (3.19),

$$\begin{aligned}
\left\| \sup_n x_n^{\frac{1}{2}} b x_n^{\frac{1}{2}} \right\|_p &= \sup \left\{ \sum_{n \geq 0} \operatorname{tr}(x_n^{\frac{1}{2}} b x_n^{\frac{1}{2}} y_n) : y_n \geq 0, \left\| \sum_{n \geq 0} y_n \right\|_{p'} \leq 1 \right\} \\
&= \sup \left\{ \sum_{n \geq 0} \operatorname{tr}(b x_n^{\frac{1}{2}} y_n x_n^{\frac{1}{2}}) : y_n \geq 0, \left\| \sum_{n \geq 0} y_n \right\|_{p'} \leq 1 \right\} \\
&\leq \|b\|_\infty \sup_n \|x_n\|_p,
\end{aligned}$$

wherein we have used the fact that

$$\operatorname{tr}(b x_n^{\frac{1}{2}} y_n x_n^{\frac{1}{2}}) \leq \|b\|_\infty \operatorname{tr}(x_n^{\frac{1}{2}} y_n x_n^{\frac{1}{2}}) = \|b\|_\infty \operatorname{tr}(x_n y_n).$$

□

The following lemma, which can be proved in the same way as in the tracial case (see [26]), will enable us to use interpolation for these norms.

Lemma 3.2. *For $2 \leq p \leq \infty$, let $\frac{1}{q} + \frac{2}{p} = 1$. Then*

$$\|(x_n)\|_{L_p^c(\mathcal{N}; \ell_\infty)} = \sup \left\{ \left(\sum_{n \geq 0} \|x_n v_n\|_2^2 \right)^{\frac{1}{2}} : \left\| \sum_{n \geq 0} v_n v_n^* \right\|_q \leq 1 \right\}.$$

Consequently, $L_p^c(\mathcal{N}; \ell_\infty)$ is a normed space.

In the following, we will restrict our attention to the following set-up. Let $\mathcal{N}$ be a σ -finite von Neumann algebra and ϕ a distinguished normal, faithful state on $\mathcal{N}$. Let $\mathcal{M} = \mathcal{N} \otimes \mathcal{B}(l_2)$ and let Tr denote the usual trace on $\mathcal{B}(l_2)$. Then $\phi \otimes \operatorname{Tr}$ is a n.s.f. weight on $\mathcal{M}$. For $1 \leq p \leq \infty$, we consider the Haagerup L_p -spaces $L_p(\mathcal{M})$ associated with this n.s.f. weight.

Let V be the space of all infinite matrices $[x_{i,j}]_{1 \leq i,j \leq \infty}$ with entries in $L_p(\mathcal{N})$, equipped with the topology of pointwise convergence in $L_p(\mathcal{N})$. More precisely, a sequence of matrices $[x_{i,j}^{(m)}]$

converges to a matrix $[x_{i,j}]$ if and only if

$$L_p(\mathcal{N}) - \lim_{m \rightarrow \infty} x_{i,j}^{(m)} = x_{i,j},$$

for all $1 \leq i, j \leq \infty$. Then V is a topological vector space. For $1 \leq p < q \leq \infty$ and $0 \leq \eta \leq 1$ define a mapping

$$\hat{I}_{q,p}^\eta = I_{q,p}^\eta \otimes \text{Id}_{\mathcal{B}(l_2)}.$$

Note that $(\hat{I}_{q,p}^\eta(L_q(\mathcal{M})), L_p(\mathcal{M}))$ forms an interpolation couple, since both embed continuously into V .

Corollary 3.3. *Let $1 \leq p < s < q \leq \infty$, $0 < \theta < 1$ such that $\frac{1}{s} = \frac{1-\theta}{p} + \frac{\theta}{q}$ and $0 \leq \eta \leq 1$. Then, completely isometrically,*

$$(3.20) \quad [\hat{I}_{q,p}^\eta(L_q(\mathcal{M})), L_p(\mathcal{M})]_\theta = \hat{I}_{s,p}^\eta(L_s(\mathcal{M})).$$

Proof. By [15], the following complete isometries hold

$$L_p(\mathcal{M}) = S_p[L_p(\mathcal{N})] \quad \text{respectively,} \quad L_q(\mathcal{M}) = S_q[L_q(\mathcal{N})].$$

Define $A_0 = I_{q,p}^\eta(L_q(\mathcal{N}))$ and $A_1 = L_p(\mathcal{N})$. By the reiteration result (3.18) it follows that $A_\theta = I_{s,p}^\eta(L_s(\mathcal{N}))$, completely isometrically. Furthermore, by [31, Corollary 1.4], we obtain a complete isometry

$$(3.21) \quad [S_q[A_0], S_p[A_1]]_\theta = S_s[A_\theta].$$

Note that for $2 \leq v < \infty$, we have

$$\| [x_{i,j} D^{\frac{1}{p} - \frac{1}{q}}] \|_{S_v[I_{v,p}(L_v(\mathcal{N}))]} = \| [x_{i,j}] \|_{S_v[L_v(\mathcal{N})]}.$$

Together with (3.21), this yields the assertion. $\square$

For $1 \leq p < \infty$, recall the spaces $L_p(\mathcal{N}; l_2^c)$ (respectively, their row version $L_p(\mathcal{N}; l_2^r)$), defined in [18] (see also [32] for the tracial case) as the completion of the family of finite sequences $a = (a_k)_{k \geq 0}$ in $L_p(\mathcal{N})$ under the norm

$$\|a\|_{L_p(\mathcal{N}; l_2^c)} = \left\| \left(\sum_{k \geq 0} |a_k|^2 \right)^{\frac{1}{2}} \right\|_p \quad \text{respectively,} \quad \|a\|_{L_p(\mathcal{N}; l_2^r)} = \left\| \left(\sum_{k \geq 0} |a_k^*|^2 \right)^{\frac{1}{2}} \right\|_p.$$

In particular, for every positive integer n , we consider the finite dimensional versions

$$\|a\|_{L_p(\mathcal{N}; (l_2^n)^c)} = \left\| \left(\sum_{k=0}^n |a_k|^2 \right)^{\frac{1}{2}} \right\|_p \quad \text{respectively,} \quad \|a\|_{L_p(\mathcal{N}; (l_2^n)^r)} = \left\| \left(\sum_{k=0}^n |a_k^*|^2 \right)^{\frac{1}{2}} \right\|_p.$$

From the above discussion, it is clear how to view the spaces $(\hat{I}_{q,p}^\eta(L_q(\mathcal{N}; \ell_2^c)), L_p(\mathcal{N}; \ell_2^c))$, where $2 \leq p < q \leq \infty$ as a compatible couple and we obtain the following:

Corollary 3.4. *Let $1 \leq p < s < q \leq \infty$, $0 < \theta < 1$ such that $\frac{1}{s} = \frac{1-\theta}{p} + \frac{\theta}{q}$ and $0 \leq \eta \leq 1$. Then, completely isometrically,*

$$(3.22) \quad [\hat{I}_{q,p}^\eta(L_q(\mathcal{N}; \ell_2^c), L_p(\mathcal{N}; \ell_2^c))_\theta = \hat{I}_{s,p}^\eta(L_s(\mathcal{N}; \ell_2^c)),$$

$$(3.23) \quad [\hat{I}_{q,p}^\eta(L_q(\mathcal{N}; \ell_2^r), L_p(\mathcal{N}; \ell_2^r))_\theta = \hat{I}_{s,p}^\eta(L_s(\mathcal{N}; \ell_2^r)),$$

Proof. Note that $L_p(\mathcal{N}; \ell_2^c)$, respectively $L_p(\mathcal{N}; \ell_2^r)$ are 1-complemented subspaces of $L_p(\mathcal{M})$ (see [18] for details). Therefore the conclusion follows from Corollary 3.3. $\square$

Remark 3.5. Corollary 3.3 holds true, more generally, in the setting of a von Neumann algebra $\mathcal{M}$ equipped with a n.f. strictly semifinite weight ψ . Recall that a n.s.f. weight ψ is called *strictly semifinite*, if there exists an increasing sequence of projections $(p_n)_{n \geq 0}$ such that $\sigma_t^\psi(p_n) = p_n$ and $\psi(p_n) < \infty$, for all $n \geq 0$.

We now establish additional properties of the spaces $L_p^c(\mathcal{M}; l_\infty)$. Let W be the space of infinite sequences of matrices $[x_{i,j}^{(n)}]_{1 \leq i,j,n \leq \infty}$ in $L_p(\mathcal{M})$. If $2 \leq p < q \leq \infty$, then both $L_p^c(\mathcal{M}; l_\infty)$ and $L_q^c(\mathcal{M}; l_\infty)$ embed continuously into the topological vector space W . In fact, the map $\hat{I}_{q,p} : L_q^c(\mathcal{M}; l_\infty) \rightarrow L_p^c(\mathcal{M}; l_\infty)$ defined by

$$\hat{I}_{q,p}([x_{i,j}^{(n)}]) = [x_{i,j}^{(n)} D^{\frac{1}{p} - \frac{1}{q}}]$$

is a continuous inclusion from $L_q^c(\mathcal{M}; l_\infty)$ into $L_p^c(\mathcal{M}; l_\infty)$ and therefore we can interpolate between these spaces.

The following result is probably well-known in interpolation theory. We include a proof for the convenience of the reader.

Lemma 3.6. *Let (A_0, A_1) be an interpolation couple with intersection Δ . Let $T_n : \Delta \rightarrow \Delta$ be a sequence of linear operators such that*

- 1) *Each T_n extends to a bounded linear operator from A_0 to A_0 such that $C_1 = \sup_n \|T_n\| < \infty$, and*
- 2) *Each T_n extends to a bounded linear operator from A_1 to A_1 such that $A_1 - \lim_{n \rightarrow \infty} T_n(x) = x$, for all $x \in \Delta$.*

Then, for all $0 < \theta < 1$ and all $x \in A_\theta = [A_0, A_1]_\theta$, we have

$$(3.24) \quad \lim_{n \rightarrow \infty} T_n(x) = x.$$

Proof. By density [3, Lemma 4.3.2] it suffices to prove (3.24) for elements $x \in A_\theta$ of the form

$$x = \sum_{k=0}^m g_k(\theta) y_k,$$

where m is a non-negative integer, g_k are analytic functions which vanish at infinity and $y_k \in \Delta$. Let $\varepsilon > 0$ and choose $n_0 \geq 0$ such that for all $n \geq n_0$, we have

$$((1 + C_1) \sum_{k=0}^m \sup_t |g_k(it)| \|y_k\|_{A_0})^{1-\theta} (\sum_{k=0}^m \sup_t |g_k(1+it)| \|T_n(y_k) - y_k\|_{A_1})^\theta < \varepsilon.$$

Since the complex interpolation method is an exact functor of exponent θ , we deduce that

$$\begin{aligned} \|x - T_n(x)\|_\theta &\leq \sup_{t \in \mathbb{R}} \left\| \sum_{k=1}^m g_k(it)(y_k - T_n(y_k)) \right\|_{A_0}^{1-\theta} \sup_{t \in \mathbb{R}} \left\| \sum_{k=1}^m g_k(1+it)(y_k - T_n(y_k)) \right\|_{A_1}^\theta \\ &\leq ((1 + C_1) \sum_{k=0}^m \sup_t |g_k(it)| \|y_k\|_{A_0})^{1-\theta} (\sum_{k=0}^m \sup_t |g_k(1+it)| \|y_k - T_n(y_k)\|_{A_1})^\theta \\ &< \varepsilon. \end{aligned}$$

This proves the assertion. $\square$

Remark 3.7. Let $2 \leq p < q \leq \infty$ and $e \in \mathcal{M}$ be a projection such that $(\phi \otimes \text{Tr})(e) < \infty$. Then $e\mathcal{M}e$ is a finite von Neumann algebra, and a similar argument as in the tracial case (see [26]) shows that the following contractive inclusion holds

$$(3.25) \quad \widehat{I}_{q,p}(L_q^c(e\mathcal{M}e; l_\infty)) \subseteq L_p^c(e\mathcal{M}e; l_\infty).$$

Proposition 3.8. *If $2 \leq p < s < q \leq \infty$ and $0 < \theta < 1$ such that $\frac{1}{s} = \frac{1-\theta}{q} + \frac{\theta}{p}$, then, isometrically,*

$$(3.26) \quad [\widehat{I}_{q,p}(L_q^c(\mathcal{M}; \ell_\infty), L_p^c(\mathcal{M}; \ell_\infty))_\theta = \widehat{I}_{s,p}(L_s^c(\mathcal{M}; \ell_\infty)).$$

Proof. We first show that we have a contractive inclusion

$$(3.27) \quad [\widehat{I}_{q,p}(L_q^c(\mathcal{M}; \ell_\infty), L_p^c(\mathcal{M}; \ell_\infty))_\theta \subseteq \widehat{I}_{s,p}(L_s^c(\mathcal{M}; \ell_\infty)).$$

Note that for each positive integer n , $L_p^c(\mathcal{M}; l_\infty^n)$ is isomorphic (as a vector space) to the n -fold tensor product $L_p(\mathcal{M}) \otimes \dots \otimes L_p(\mathcal{M})$ (n times). Therefore $(\widehat{I}_{q,p}^{(n)}(L_q^c(\mathcal{M}; l_\infty^n), L_p^c(\mathcal{M}; l_\infty^n)))$ is a compatible couple, where

$$\widehat{I}_{q,p}^{(n)} = \widehat{I}_{q,p} \otimes \text{Id}(l_\infty^n).$$

By Remark 3.1 (1), it suffices to show that for all $n \geq 1$ we have, contractively,

$$(3.28) \quad [\widehat{I}_{q,p}^{(n)}(L_q^c(\mathcal{M}; l_\infty^n), L_p^c(\mathcal{M}; l_\infty^n))_\theta \subseteq \widehat{I}_{s,p}^{(n)}(L_s^c(\mathcal{M}; l_\infty^n)).$$

In the following we fix a positive integer n and consider only sequences of length n . We claim that it suffices to prove that for all $m \geq 1$ the following contractive inclusion holds

$$(3.29) \quad [\widehat{I}_{q,p}^{(n)}(L_q^c(\mathcal{N} \otimes M_m; l_\infty^n), L_p^c(\mathcal{N} \otimes M_m; l_\infty^n))_\theta \subseteq \widehat{I}_{q,p}^{(n)}(L_s^c(\mathcal{N} \otimes M_m; l_\infty^n)),$$

where M_m is the algebra of $m \times m$ complex matrices. Note that $\mathcal{N} \otimes M_m = e_m \mathcal{M} e_m$, where $e_m = 1 \otimes p_m$ and $p_m : \mathcal{B}(l_2) \rightarrow M_m$ is the canonical orthogonal projection. Set

$$A_0 = \widehat{I}_{q,p}^{(n)}(L_q^c(\mathcal{N} \otimes M_m; l_\infty^n)) \quad \text{and} \quad A_1 = L_p^c(\mathcal{N} \otimes M_m; l_\infty^n).$$

By Remark 3.7, their intersection and sum are, respectively,

$$\Delta_0 = A_0 \cap A_1 = \widehat{I}_{q,p}^{(n)}(L_q^c(\mathcal{N} \otimes M_m; l_\infty^n)), \quad \Sigma_0 = A_0 + A_1 = L_p^c(\mathcal{N} \otimes M_m; l_\infty^n).$$

Consider the map $S_m : L_q^c(\mathcal{M}; l_\infty^n) \rightarrow L_q^c(e_m \mathcal{M} e_m; l_\infty^n)$ defined by

$$S_m((x_k)_{k=1}^n) = (e_m x_k e_m)_{k=1}^n.$$

By (2) in Remark 3.1, it follows that

$$\sup_m \|S_m\|_{A_0 \rightarrow A_0} \leq 1.$$

Furthermore, we have $A_1 - \lim_{m \rightarrow \infty} S_m(x) = x$, for all $x \in \Delta_0$ (see [14]). An application of Lemma 3.6 shows that, for all $x \in [A_0, A_1]_\theta$,

$$\lim_{m \rightarrow \infty} S_m(x) = x.$$

Therefore the claim is justified. We now prove (3.29). Let $2 \leq v < \infty$ such that $\frac{1}{v} = \frac{1}{p} - \frac{1}{q}$. Furthermore, define $f(v)$ by the relation $\frac{1}{f(v)} + \frac{1}{v} = \frac{1}{2}$ and note that $\frac{1}{f(q)} - \frac{1}{f(p)} = \frac{1}{v}$. Let

$$B_0 = \widehat{I}_{f(p), f(q)}^{(n)}(L_{f(p)}(\mathcal{N} \otimes M_m; (l_2^n)^r)), \quad B_1 = \widehat{I}_{f(q), f(q)}^{(n)}(L_{f(q)}(\mathcal{N} \otimes M_m; (l_2^n)^r)).$$

Then, their intersection and sum are, respectively,

$$\Delta_1 = \widehat{I}_{f(p), f(q)}^{(n)}(L_{f(p)}(\mathcal{N} \otimes M_m; (l_2^n)^r)), \quad \Sigma_1 = \widehat{I}_{f(q), f(q)}^{(n)}(L_{f(q)}(\mathcal{N} \otimes M_m; (l_2^n)^r)).$$

Furthermore, given finite sequences of $m \times m$ matrices $([x_{i,j}^{(k)} D_v^{\frac{1}{v}}]_{1 \leq i, j \leq m})_{k=1}^n$ in Δ_0 and, respectively, $([D_v^{\frac{1}{v}} y_{i,j}^{(k)}]_{1 \leq i, j \leq m})_{k=1}^n$ in Δ_1 , define

$$T\left(\left([x_{i,j}^{(k)} D_v^{\frac{1}{v}}]_k\right) \times \left([D_v^{\frac{1}{v}} y_{i,j}^{(k)}]_k\right)\right) = \left([x_{i,j}^{(k)} D_v^{\frac{1}{v}} y_{i,j}^{(k)}]_k\right) \in \ell_2^n(L_2(\mathcal{N} \otimes M_m)).$$

This map is well-defined, since for each $1 \leq k \leq n$, $[x_{i,j}^{(k)}]_{1 \leq i, j \leq m} \in L_q(\mathcal{N} \otimes M_m)$ and $[y_{i,j}^{(k)}]_{1 \leq i, j \leq m} \in L_{f(p)}(\mathcal{N} \otimes M_m)$, and, by construction, $\frac{1}{q} + \frac{1}{v} + \frac{1}{f(q)} = \frac{1}{2}$. Moreover, we deduce from Lemma 3.2 that

$$\|T : \Delta_0 \times \Delta_1 \rightarrow \ell_2^n(L_2(\mathcal{N} \otimes M_m))\| \leq 1.$$

Using multilinear complex interpolation (see [3, Theorem 4.4.1]), the map T extends to a contraction

$$T : [A_0, A_1]_\theta \times [B_0, B_1]_\theta \rightarrow \ell_2^n(L_2(\mathcal{N} \otimes M_m)).$$

By Corollary 3.4 applied to the algebra $\mathcal{N} \otimes M_m$, we deduce that, isometrically,

$$[B_0, B_1]_\theta = \widehat{I}_{f(s), f(q)}^{(n)}(L_{f(s)}(\mathcal{N} \otimes M_m; (l_2^n)^r)).$$

Therefore the extended map

$$T : [A_0, A_1]_\theta \times \widehat{I}_{f(s), f(q)}^{(n)}(L_{f(s)}(\mathcal{N} \otimes M_m; (l_2^n)^r)) \rightarrow \ell_2^n(L_2(\mathcal{N} \otimes M_m))$$

is a contraction. A further application of Lemma 3.2 yields the estimate

$$\left\| T : [\widehat{I}_{q,p}^{(n)}(L_q^c(\mathcal{N} \otimes M_m; \ell_\infty^n)), L_p^c(\mathcal{N} \otimes M_m; \ell_\infty^n)]_\theta \rightarrow \widehat{I}_{s,p}^{(n)}(L_s^c(\mathcal{N} \otimes M_m; \ell_\infty^n)) \right\| \leq 1.$$

Thus (3.29) is justified. The proof of the reverse inclusion in (3.26) reduces, just as in the tracial case, to the interpolation result (3.20), via the factorization property of the spaces $L_p^c(\mathcal{M}; l_\infty)$, proved in [14]. $\square$

Let us now turn our attention to martingales invariant under the automorphism group of a given state. Indeed, we will assume that $(\mathcal{N}_n)_{n \geq 0}$ is an increasing sequence of von Neumann subalgebras of $\mathcal{N}$ such that $\sigma_t(\mathcal{N}_n) \subset \mathcal{N}_n$ for all $t \in \mathbb{R}$ and all $n \geq 0$. Moreover, let $(p_n)_{n \geq 0}$ be the increasing sequence of projections given by the units in $\mathcal{N}_n$. We will further assume that $\sigma_t(p_n) = p_n$ for all $t \in \mathbb{R}$ and all $n \geq 0$. For fixed $n \in \mathbb{N}$, according to [42], there exists a conditional expectation $\tilde{\mathcal{E}}_n : p_n \mathcal{N} p_n \rightarrow \mathcal{N}_n$ such that

$$(3.30) \quad \tilde{\mathcal{E}}_n(\sigma_t(x)) = \sigma_t(\tilde{\mathcal{E}}_n(x))$$

for all $x \in p_n \mathcal{N} p_n$ and all $t \in \mathbb{R}$. Let us define $\mathcal{E}_n(x) = \tilde{\mathcal{E}}_n(p_n x p_n)$ and note that $\mathcal{E}_n$ still satisfies the invariance property (3.30). Then the martingale differences are defined by

$$d_n(x) = \mathcal{E}_n(x) - \mathcal{E}_{n-1}(x),$$

where $\mathcal{E}_{-1}(x) = 0$. Moreover, for $1 \leq p < \infty$, the conditional expectation and the martingale difference operators extend in a natural way to $L_p(\mathcal{N})$ (see [18]), such that for all $0 < \theta < 1$ the relation

$$(3.31) \quad \mathcal{E}_n(D^{\frac{1-\theta}{p}} x D^{\frac{\theta}{p}}) = D^{\frac{1-\theta}{p}} \mathcal{E}_n(x) D^{\frac{\theta}{p}}$$

holds for all $x \in \mathcal{N}$. Furthermore, if $\frac{1}{p} = \frac{1}{q} + \frac{1}{s} + \frac{1}{t}$, then for all $a \in L_s(\mathcal{N}_n)$, $b \in L_t(\mathcal{N}_n)$ and $x \in L_q(\mathcal{N})$, we have

$$(3.32) \quad \mathcal{E}_n(axb) = a \mathcal{E}_n(x) b.$$

Recall also that if $1 \leq p \leq \infty$ and $x_\infty \in L_p(\mathcal{N})$, then the sequence $(x_n)_{n \geq 0}$ defined by $x_n = \mathcal{E}_n(x_\infty)$ is a bounded $L_p(\mathcal{N})$ -martingale which converges to x_∞ in $L_p(\mathcal{N})$ (respectively, in the w^* -topology if $p = \infty$). Conversely, for $1 < p < \infty$, we deduce from the uniform convexity of $L_p(\mathcal{N})$ that every bounded $L_p(\mathcal{N})$ -martingale converges to some element $x_\infty \in L_p(\mathcal{N})$ and hence it is of the above form. Therefore, we will often identify a martingale with its limit, whenever this exists.

For convenience, we will also assume that the predual $\mathcal{N}_*$ is separable. Our results will hold without this assumption and we refer to the Appendix in [12] for techniques deducing the general case from the σ -finite one. However, assuming that $\mathcal{N}_*$ is separable, we will find a separable σ -weakly dense subalgebra $\mathcal{A}_0$ of $\mathcal{N}_a$. Following [14], we can then consider the countably generated

Hilbert C^* -module F_n generated by $\mathcal{N}_n$ and $\mathcal{A}_0$, and find a right module isomorphism $u_n : F_n \rightarrow C_\infty(\mathcal{N}_n)$ such that for all $x, y \in F_n$, we have

$$(3.33) \quad \mathcal{E}_n(y^*x) = u_n(y)^*u_n(x).$$

Moreover, for $1 \leq p < \infty$ the map $u_n^p : \mathcal{N}_{n,a}D^{\frac{1}{p}} \rightarrow C_\infty(\mathcal{N}_{n,a})D^{\frac{1}{p}}$ defined by

$$u_n^p(xD^{\frac{1}{p}}) = u_n(x)D^{\frac{1}{p}}, \quad x \in \mathcal{N}_{n,a}$$

is well-defined and isometric if we equip $\mathcal{N}_{n,a}D^{\frac{1}{p}}$ with the norm

$$\left\| xD^{\frac{1}{p}} \right\|_{L_p(\mathcal{N}, \mathcal{E}_n)} = \left\| D^{\frac{1}{p}} \mathcal{E}_n(x^*x) D^{\frac{1}{p}} \right\|_{\frac{p}{2}}^{\frac{1}{2}}.$$

Indeed, in [14] this was considered only in the faithful case. However, by working with the algebra $\mathcal{N}_n + \mathbb{C}(1 - p_n)$ and the corresponding $\tilde{u}_n^p$, and then defining $u_n^p(xD^{\frac{1}{p}}) = \tilde{u}_n^p(xD^{\frac{1}{p}})p_n$, we can extend the construction to the non-faithful case (see Section 8 in [18] for details).

For $2 \leq p \leq \infty$, we will consider the space $L_p^c MO(\mathcal{N})$ of martingale difference sequences $x \cong (d_k)_{k \geq 0}$ satisfying

$$(3.34) \quad \|x\|_{L_p^c MO(\mathcal{N})} = \sup_m \left\| \sup_{n \leq m} \mathcal{E}_n \left(\sum_{k=n}^m d_k^* d_k \right) \right\|_{\frac{p}{2}}^{\frac{1}{2}} < \infty.$$

Note that if $p = \infty$, then $L_\infty^c MO(\mathcal{N}) = BMO_c(\mathcal{N})$.

Recall also the following norms

$$(3.35) \quad \|x\|_{H_p^c(\mathcal{N})} = \|dx\|_{L_p(\mathcal{N}; l_2^c)}.$$

The next results concerning the spaces $L_p^c MO(\mathcal{N})$ will be very useful in the sequel. The following noncommutative version of Stein's inequality is essentially contained in [18].

Lemma 3.9. *Let $1 < p < \infty$. Define a map Q on all finite sequences $x = (x_n)_{n \geq 0}$ in $L_p(\mathcal{N})$ by $Q(x) = (\mathcal{E}_n(x_n))_{n \geq 0}$. Then*

$$\|Q(x)\|_{L_p(\mathcal{N}; l_2^c)} \leq \gamma_p \|x\|_{L_p(\mathcal{N}; l_2^c)} \quad \text{respectively,} \quad \|Q(x)\|_{L_p(\mathcal{N}; l_2^r)} \leq \gamma_p \|x\|_{L_p(\mathcal{N}; l_2^r)}.$$

Thus Q extends to a bounded projection on $L_p(\mathcal{N}; l_2^c)$ and $L_p(\mathcal{N}; l_2^r)$, respectively.

Furthermore, as an application of Stein's inequality, Doob's inequality and the duality between $L_p^c MO(\mathcal{N})$ and $H_{p'}^c(\mathcal{N})$, where $\frac{1}{p} + \frac{1}{p'} = 1$, the following was proved in [14] (see also Section 8 in [18]):

Theorem 3.10. *Let $1 < p < \infty$, then, with equivalent norms,*

$$H_p^c(\mathcal{N}) = L_p^c MO(\mathcal{N}).$$

Let $2 \leq p < q \leq \infty$. As already observed in [18], the map $I_{q,p} : L_q^c MO(\mathcal{N}) \rightarrow L_p^c MO(\mathcal{N})$ is a contraction. However, for $0 < \eta < 1$, the inclusion map $I_{q,p}^\eta : L_q^c MO(\mathcal{N}) \rightarrow L_p^c MO(\mathcal{N})$ is not necessarily continuous anymore, but well-defined on finite sequences. Since we would like to consider interpolation between these spaces, let $\mathcal{S} = \prod_{n \in \mathbb{N}} \prod_{m \leq n} L_p(\mathcal{N}_m)$, endowed with the product topology. For $p < v < q$ with $\frac{1}{p} = \frac{1}{q} + \frac{1}{v}$, we may consider the embedding $\tilde{I}_{q,p}^\eta : L_q^c MO(\mathcal{N}) \rightarrow \mathcal{S}$ defined by

$$(3.36) \quad \tilde{I}_{q,p}^\eta((x_m)_m) = (D^{\frac{1-\eta}{v}}(x_m - x_{n-1})D^{\frac{\eta}{v}})_{m,n}$$

where $(x_m)_m$ is a finite martingale. Note that by theorem 3.10, it follows that finite martingales are dense in $L_q^c MO(\mathcal{N})$ for $q < \infty$, since this is true for the space $H_q^c(\mathcal{N})$.

The following proposition is a non-tracial modification of the result in [26].

Proposition 3.11. *Let $2 \leq p < s < q \leq \infty$ and $0 \leq \theta \leq 1$ such that $\frac{1}{s} = \frac{1-\theta}{q} + \frac{\theta}{p}$, then, with equivalent norms*

$$(3.37) \quad [\tilde{I}_{q,p}(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})]_\theta = \tilde{I}_{s,p}(L_s^c MO(\mathcal{N})).$$

Proof. For $2 \leq v \leq \infty$, we define a map $\Phi_v : L_v^c MO(\mathcal{N}) \rightarrow L_v^c(\mathcal{M}; \ell_\infty)$ by

$$\Phi_v(x) = (u_n^v(x - \mathcal{E}_{n-1}(x)))_{n \geq 0}.$$

We claim that Φ_v is an isometry. Indeed, an application of Lemma 3.2, together with (3.33) shows that

$$\begin{aligned} \|\Phi_v(x)\|_{L_v^c(\mathcal{M}; \ell_\infty)} &= \left\| \sup_n u_n^v(x - \mathcal{E}_{n-1}(x))^* u_n^v(x - \mathcal{E}_{n-1}(x)) \right\|_{\frac{v}{2}}^{\frac{1}{2}} \\ &= \left\| \sup_n \mathcal{E}_n((x - \mathcal{E}_{n-1}(x))^*(x - \mathcal{E}_{n-1}(x))) \right\|_{\frac{v}{2}}^{\frac{1}{2}} \\ &= \|x\|_{L_v^c MO(\mathcal{N})}. \end{aligned}$$

Moreover, we observe that by definition of the maps u_n^v we have

$$\Phi_p(x D^{\frac{1}{p} - \frac{1}{s}}) = \Phi_s(x) D^{\frac{1}{p} - \frac{1}{s}}.$$

This implies that

$$\Phi_p \tilde{I}_{s,p} = \widehat{I}_{s,p} \Phi_s.$$

Hence, the family Φ_v is compatible with the interpolation couples

$$(\tilde{I}_{q,p}(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})) \quad \text{and} \quad (\widehat{I}_{q,p}(L_q^c(\mathcal{M}, \ell_\infty)), L_p^c(\mathcal{M}, \ell_\infty)).$$

We deduce that

$$\Phi_p : [\tilde{I}_{q,p}(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})]_\theta \rightarrow [\widehat{I}_{q,p}(L_q^c(\mathcal{M}; \ell_\infty)), L_p^c(\mathcal{M}; \ell_\infty)]_\theta$$

is a contraction. An application of Proposition 3.8 yields the contraction

$$\widehat{I}_{s,p}\Phi_p : [\tilde{I}_{q,p}(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})]_\theta \rightarrow \widehat{I}_{s,p}(L_s^c(\mathcal{M}; \ell_\infty)) .$$

By considering elements in the intersection $\Delta = (L_q^c MO(\mathcal{N}))D^{\frac{1}{p}-\frac{1}{q}} \subset L_p^c MO(\mathcal{N})$, we deduce that the image of $\widehat{I}_{s,p}\Phi_p$ is contained in $\widehat{I}_{s,p}\Phi_s(L_s^c MO(\mathcal{N}))$. Since Φ_s is isometric, we obtain a contractive inclusion

$$(3.38) \quad j_s : [\tilde{I}_{q,p}(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})]_\theta \rightarrow \tilde{I}_{s,p}(L_s^c MO(\mathcal{N})) ,$$

densely defined by the formula

$$j_s(xD^{\frac{1}{p}-\frac{1}{q}}) = xD^{\frac{1}{p}-\frac{1}{s}} .$$

A similar argument as in the proof of Proposition 3.4 in [26] yields the converse inclusion. Namely, since the unit ball of the interpolation space $[\tilde{I}_{q,p}(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})]_\theta$ is dense in the unit ball of the space $[\tilde{I}_{q,p}(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})]^\theta$ with respect to the sum topology, we also obtain the contractive inclusion $[\tilde{I}_{q,p}(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})]^\theta \subseteq \tilde{I}_{s,p}(L_s^c MO(\mathcal{N}))$. Therefore, to show the converse inclusion in (3.38) it follows by duality that it suffices to prove

$$(3.39) \quad [H_{q'}^c(\mathcal{N}), \tilde{I}_{p',q'}(H_{p'}^c(\mathcal{N}))]_\theta \subseteq \tilde{I}_{s',q'}(H_{s'}^c(\mathcal{N})) ,$$

where p', q', s' are respectively the conjugates of p, q, s . Note that since $\frac{1}{s} = \frac{1-\theta}{q} + \frac{\theta}{p}$, we deduce that $1 \leq q' < s' < p' \leq 2$ and $\frac{1}{s'} = \frac{1-\theta}{q'} + \frac{\theta}{p'}$. Therefore, by Corollary 3.4 we obtain

$$(3.40) \quad [H_{q'}^c(\mathcal{N}), \tilde{I}_{p',q'}(H_{p'}^c(\mathcal{N}))]_\theta \subseteq [L_{q'}^c(\mathcal{N}; l_2^c), \hat{I}_{p',q'}(L_{p'}^c(\mathcal{N}; l_2^c))]_\theta = \hat{I}_{s',q'}(L_{s'}^c(\mathcal{N}; l_2^c)) .$$

By Stein's inequality (see Lemma 3.9) we have for all $x = (x_m)_m \in H_{q'}^c(\mathcal{N}) \subseteq \tilde{I}_{s',q'}(H_{s'}^c(\mathcal{N}))$,

$$\left\| \left(\sum d_m(x_m)^* d_m(x_m) \right)^{\frac{1}{2}} \right\|_{s'} \leq \gamma_{s'} \left\| \left(\sum x_m^* x_m \right)^{\frac{1}{2}} \right\|_{s'} ,$$

where the constant $\gamma_{s'}$ depends only on s' . Together with (3.40), this implies (3.39) and the assertion is proved. $\square$

As an application, we prove the result announced in [18].

Corollary 3.12. *Let $4 < p < \infty$, then for every $x \in L_p(\mathcal{N})$, we have*

$$\|x\|_{L_p^c MO(\mathcal{N})} \leq c_4^{4/p} \|x\|_p .$$

Here c_4 is an absolute constant.

Proof. Fix $0 < \theta < 1$ such that $\frac{1}{p} = \frac{1-\theta}{\infty} + \frac{\theta}{4}$. Let $x \in L_p(\mathcal{N})$ with $\|x\|_p \leq 1$. Consider the polar decomposition $x = u|x|$ with $u \in \mathcal{N}$ and $|x| = (x^*x)^{\frac{1}{2}} \in L_p(\mathcal{N})$. It follows that $\|x\|_p = \|x\|_p \leq 1$. Given $\delta > 0$ and $C \geq 1$ define a function

$$f(z) = C^{-z} \exp(\delta(\theta - z)^2) u|x|^{\frac{zp}{4}} D^{\frac{(1-z)}{4}} ,$$

for all z in the unit strip. According to Fact 2.1 in the preliminaries, it follows that f is analytic and $f(z) \in L_4(\mathcal{N})$. Furthermore, for all $t \in \mathbb{R}$, note that $|x|^{pit} D^{-it} \in \mathcal{N}$. Hence

$$(3.41) \quad \left\| f(it) D^{-\frac{1}{4}} \right\|_{\infty} \leq \exp(\delta \theta^2),$$

and, respectively,

$$(3.42) \quad \|f(1+it)\|_4 \leq C^{-1} \exp(\delta \theta^2).$$

According to Theorem 3.10 (see [18] for details), there exist absolute constants $\beta_4, \gamma_4 > 0$ such that for all $y \in L_4(\mathcal{N})$,

$$\|y\|_{L_4^c MO(\mathcal{N})} \leq 2\gamma_4 \|y\|_{H_4^c(\mathcal{N})} \leq 2\gamma_4 \beta_4 \|y\|_4.$$

Trivially, we have for $y \in \mathcal{N}$ that

$$\|y\|_{BMO_c(\mathcal{N})} \leq 2 \|y\|_{\infty}.$$

Choose $C = 2\gamma_4 \beta_4 := c_4$. It follows that $f(\theta) \in [\tilde{I}_{\infty,4}(BMO_c(\mathcal{N})), L_4^c MO(\mathcal{N})]_{\theta}$, with

$$\|f(\theta)\|_{\theta} \leq (\exp(\delta \theta^2))^{1-\theta} (\exp(\delta \theta^2))^{\theta} = \exp(\delta).$$

According to Proposition 3.11, we deduce that

$$\left\| \tilde{I}_{p,4}(f(\theta)) \right\|_{L_p^c MO(\mathcal{N})} \leq \exp(\delta).$$

Since $\delta > 0$ was arbitrarily chosen and $f(\theta) = c_4^{-\frac{4}{p}} x D^{\frac{1}{4} - \frac{1}{p}}$, the assertion follows. $\square$

Lemma 3.13. *Let $2 \leq p < q \leq \infty$, $0 < \theta < 1$ and $0 \leq \eta \leq 1$. Then $\bigcup_{m \geq 0} \tilde{I}_{q,p}((\mathcal{N}_{m,a}) D^{\frac{1}{q}})$ is norm dense in the interpolation space $[\tilde{I}_{q,p}^{\eta}(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})]_{\theta}$.*

Proof. First we note that the sequence $(\mathcal{E}_m)_{m \geq 0}$ given by the conditional expectations satisfies the assumptions of Lemma 3.6. Thus the space of finite martingales is norm dense in the interpolation space, since this is true for the space $L_p^c MO(\mathcal{N})$, as observed before. Given an integer $m \geq 0$, we note that the inclusions

$$L_s(\mathcal{N}_m; l_2^c) \subset L_s^c MO(\mathcal{N}_m) \subset L_s^c MO(\mathcal{N})$$

are continuous, since

$$\|d_k(x)\|_{L_s(\mathcal{N}_m)} \leq 2 \min\{\|x\|_{L_s^c MO(\mathcal{N}_m)}, \|x\|_{L_s(\mathcal{N}_m)}\}$$

for all $0 \leq k \leq m$. The intersection space is

$$\tilde{I}_{q,p}^{\eta}(L_q^c MO(\mathcal{N}_m)) \cap L_p^c MO(\mathcal{N}_m) = \tilde{I}_{q,p}^{\eta}(L_q^c MO(\mathcal{N}_m)).$$

Since $\mathcal{N}_m D^{\frac{1}{q}}$ is norm dense in $L_q(\mathcal{N}_m)$, we deduce that $\tilde{I}_{q,p}^\eta(\mathcal{N}_m D^{\frac{1}{q}})$ is also dense in the interpolation space. In order to obtain analytic elements we recall the approximation operators $R_k : L_p(\mathcal{N}) \rightarrow L_p(\mathcal{N})$ defined by

$$R_k(x) = k^{\frac{1}{2}} \pi^{-\frac{1}{2}} \int_{\mathbb{R}} \exp(-kt^2) D^{it} x D^{-it} dt .$$

Since $\sigma_t(x) = D^{it} x D^{-it}$ is strongly continuous on $\mathcal{N}$, we deduce by approximation that σ_t is norm continuous on $L_p(\mathcal{N})$ for all $2 \leq p < \infty$ (see [14], [17]). It then follows that

$$L_p - \lim_{k \rightarrow \infty} R_k(x) = x$$

for every $x \in L_p(\mathcal{N})$. Therefore, the same is true for the space $L_p^c MO(\mathcal{N}_m)$. Clearly, the map $\sigma_t : L_p^c MO(\mathcal{N}) \rightarrow L_p^c MO(\mathcal{N})$ is a contraction. It follows that R_k is a contraction on $L_p^c MO(\mathcal{N})$, for all $2 \leq p \leq \infty$. In view of Lemma 3.6, we deduce that elements of the form $\tilde{I}_{q,p}^\eta(R_k(x) D^{\frac{1}{q}})$ are norm dense in the interpolation space $[\tilde{I}_{q,p}^\eta(L_q^c MO(\mathcal{N}_m)), L_p^c MO(\mathcal{N}_m)]_\theta$, for all $x \in \mathcal{N}_m$. Using the fact that the image $R_k(\mathcal{N}_m)$ consists of analytic elements and

$$R_k(x D^{\frac{1}{q}}) = R_k(x) D^{\frac{1}{q}} ,$$

we deduce the assertion. $\square$

In order to establish the inclusion $I_{\infty,p}^1(BMO(\mathcal{N})) = I_{\infty,p}^1(BMO_c \cap BMO_r) \subset L_p(\mathcal{N})$, we have to consider and analyze the space $I_{\infty,p}^1(BMO_r(\mathcal{N}))$. In contrast with the inclusion $I_{\infty,p}^1(BMO_c(\mathcal{N})) \subset L_p^c MO(\mathcal{N})$, which follows easily by the definition, for the row version we are multiplying by $D^{\frac{1}{p}}$ on the wrong side. Alternatively, we may consider the space $I_{\infty,p}^0(BMO_c(\mathcal{N}))$. In this case we can use Kosaki's change of density argument (see [23]) in order to investigate the interpolation space $[I_{\infty,p}^0(BMO_c(\mathcal{N}), L_p^c MO(\mathcal{N}))]_\theta$, for $0 < \theta < 1$.

Proposition 3.14. *Let $2 \leq p < q \leq \infty$, $p \leq v < \infty$ such that $\frac{1}{p} = \frac{1}{q} + \frac{1}{v}$ and $0 < \theta < 1$. Then for all $0 \leq \eta < 1$, the spaces $[\tilde{I}_{q,p}^\eta(L_q^c MO(\mathcal{N}), L_p^c MO(\mathcal{N}))]_\theta$ and $[\tilde{I}_{q,p}^1(L_q^c MO(\mathcal{N}), L_p^c MO(\mathcal{N}))]_\theta$ are isometrically isomorphic. The isomorphism*

$$T_\theta : [\tilde{I}_{q,p}^1(L_q^c MO(\mathcal{N}), L_p^c MO(\mathcal{N}))]_\theta \rightarrow [\tilde{I}_{q,p}^\eta(L_q^c MO(\mathcal{N}), L_p^c MO(\mathcal{N}))]_\theta$$

is defined by

$$T_\theta((x_m D^{\frac{1}{v}})_m) = \left(D^{\frac{(1-\eta)(1-\theta)}{v}} x_m D^{\frac{1}{v} - \frac{(1-\eta)(1-\theta)}{v}} \right)_m ,$$

for all adapted finite sequences $(x_m)_m \in L_q^c MO(\mathcal{N})$.

Proof. We will first construct a contraction

$$T_\theta : [\tilde{I}_{q,p}^1(L_q^c MO(\mathcal{N}), L_p^c MO(\mathcal{N}))]_\theta \rightarrow [\tilde{I}_{q,p}^\eta(L_q^c MO(\mathcal{N}), L_p^c MO(\mathcal{N}))]_\theta .$$

According to Lemma 3.13, it suffices to define T_θ on the dense set $\bigcup_{m \geq 0} \tilde{I}_{q,p}^1(\mathcal{N}_{m,a} D^{\frac{1}{q}})$ and show that the restriction is contractive. Furthermore, by the density result [3, Lemma 4.3.2], it is enough to consider elements of the form

$$x = f(\theta) D^{\frac{1}{p}},$$

where

$$f(z) = \sum_{k=1}^l g_k(z) y_k$$

with $y_k \in \mathcal{N}_{m,a}$, for some m , and g_k analytic functions which vanish at infinity such that

$$\max \left\{ \sup_{t \in \mathbb{R}} \left\| f(it) D^{\frac{1}{q}} \right\|_{L_q^c MO(\mathcal{N})}, \sup_{t \in \mathbb{R}} \left\| f(1+it) D^{\frac{1}{p}} \right\|_{L_p^c MO(\mathcal{N})} \right\} < 1.$$

For all z in the unit strip S we define

$$g(z) = D^{\frac{(1-\eta)(1-z)}{v}} f(z) D^{\frac{1}{q} + \frac{\eta}{v} + \frac{(1-\eta)z}{v}} = \sum_{k=1}^l g_k(z) (D^{\frac{(1-\eta)(1-z)}{v}} y_k D^{\frac{(1-\eta)z}{v}} D^{\frac{1}{q} + \frac{\eta}{v}}).$$

According to Fact 2.1, this function is analytic, taking values in $L_p(\mathcal{N}_m) \subset L_p^c(\mathcal{N}_m; l_\infty)$. Then, we observe that σ_t is an isometry on $L_p^c MO(\mathcal{N})$ and therefore

$$\begin{aligned} \|g(1+it)\|_{L_p^c MO(\mathcal{N})} &= \left\| D^{-\frac{it(1-\eta)}{v}} f(1+it) D^{\frac{1}{q} + \frac{\eta}{v} + \frac{1-\eta}{v}} D^{\frac{(1-\eta)it}{v}} \right\|_{L_p^c MO(\mathcal{N})} \\ &= \left\| f(1+it) D^{\frac{1}{p}} \right\|_{L_p^c MO(\mathcal{N})} < 1. \end{aligned}$$

On the other hand, by (3.36),

$$\begin{aligned} \left\| D^{-\frac{1-\eta}{v}} g(it) D^{-\frac{\eta}{v}} \right\|_{L_q^c MO(\mathcal{N})} &= \left\| D^{-\frac{it(1-\eta)}{v}} f(it) D^{\frac{(1-\eta)it}{v}} D^{\frac{1}{q}} \right\|_{L_q^c MO(\mathcal{N})} \\ &= \left\| D^{-\frac{it(1-\eta)}{v}} f(it) D^{\frac{1}{q}} D^{\frac{(1-\eta)it}{v}} \right\|_{L_q^c MO(\mathcal{N})} \\ &= \left\| f(it) D^{\frac{1}{q}} \right\|_{L_q^c MO(\mathcal{N})} < 1. \end{aligned}$$

Therefore, for $x = f(\theta) D^{\frac{1}{p}}$ it follows that

$$\begin{aligned} (3.43) \quad T_\theta(x) = g(\theta) &= \sum_{k=1}^l g_k(\theta) D^{\frac{(1-\eta)(1-\theta)}{v}} y_k D^{\frac{(1-\eta)\theta}{v}} D^{\frac{1}{q} + \frac{\eta}{v}} \\ &= D^{\frac{(1-\eta)(1-\theta)}{v}} f(\theta) D^{\frac{1}{q}} D^{\frac{(1-\eta)\theta}{v} + \frac{\eta}{v}} = D^{\frac{(1-\eta)(1-\theta)}{v}} f(\theta) D^{\frac{1}{p} - \frac{(1-\eta)(1-\theta)}{v}} \end{aligned}$$

is an element in the interpolation space $[\tilde{I}_{q,p}^\eta(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})]_\theta$. Hence T_θ defined by (3.43) extends to a contraction. Conversely, a similar argument shows that the map $\tilde{T}_\theta$ defined by

$$\tilde{T}_\theta(D^{\frac{1-\eta}{v}} f(\theta) D^{\frac{1}{q}} D^{\frac{\eta}{v}}) = D^{\frac{(1-\eta)\theta}{v}} f(\theta) D^{\frac{1}{q} + \frac{\eta}{v} + \frac{(1-\eta)(1-\theta)}{v}}$$

is a contraction. Since f is analytic, there exists $y \in \mathcal{N}_{m,a}$ such that

$$D^{\frac{(1-\eta)\theta}{v}} f(\theta) = y D^{\frac{(1-\eta)\theta}{v}}.$$

Furthermore, we deduce that

$$\begin{aligned} T_\theta \tilde{T}_\theta (D^{\frac{1-\eta}{v}} f(\theta) D^{\frac{1}{q}} D^{\frac{\eta}{v}}) &= T_\theta (y D^{\frac{1}{p}}) \\ &= D^{\frac{(1-\eta)(1-\theta)}{v}} y D^{\frac{1}{p} - \frac{(1-\eta)(1-\theta)}{v}} \\ &= D^{\frac{(1-\eta)(1-\theta)}{v}} y D^{\frac{(1-\eta)\theta}{v}} D^{\frac{1}{q} + \frac{\eta}{v}} \\ &= D^{\frac{1-\eta}{v}} f(\theta) D^{\frac{1}{q} + \frac{\eta}{v}}. \end{aligned}$$

Hence $T_\theta \tilde{T}_\theta = \text{id}$. Similarly, we may show that $\tilde{T}_\theta T_\theta = \text{id}$, and the assertion follows. $\square$

Corollary 3.15. *Let $2 < p < s < q \leq \infty$, $0 < \theta < 1$ such that $\frac{1}{s} = \frac{1-\theta}{q} + \frac{\theta}{p}$, and $0 \leq \eta \leq 1$. Then, isometrically,*

$$[\tilde{I}_{q,p}^\eta(L_q^c MO(\mathcal{N})), L_p^c MO(\mathcal{N})]_\theta = \tilde{I}_{s,p}^\eta(L_s^c MO(\mathcal{N})).$$

Proof. Let us note first that $\frac{1}{p} - \frac{1}{s} = \frac{1-\theta}{v}$, where $\frac{1}{v} = \frac{1}{p} - \frac{1}{q}$. Then, we observe that

$$T_\theta(x D^{\frac{1}{p} - \frac{1}{s}}) = D^{(1-\eta)\frac{1-\theta}{v}} x D^{\frac{1}{p} - \frac{1}{s} - (1-\eta)\frac{1-\theta}{v}} = D^{(1-\eta)(\frac{1}{p} - \frac{1}{s})} x D^{\eta(\frac{1}{p} - \frac{1}{s})}.$$

The assertion follows now from Proposition 3.11 and Proposition 3.14. $\square$

Recall now the row version of the martingale norm (3.34)

$$\|x\|_{L_p^r MO(\mathcal{N})} = \sup_m \left\| \sup_{n \leq m} \mathcal{E}_n \left(\sum_{k=n}^m d_k(x) d_k(x)^* \right) \right\|_{\frac{p}{2}}^{\frac{1}{2}}.$$

Interchanging the roles of columns and rows we immediately get the following result.

Proposition 3.16. *Let $2 < p < t < q \leq \infty$, $0 < \theta < 1$ such that $\frac{1}{s} = \frac{1-\theta}{q} + \frac{\theta}{p}$, and $0 \leq \eta \leq 1$. Then, isometrically,*

$$[\tilde{I}_{q,p}^\eta(L_q^r MO(\mathcal{N})), L_p^r MO(\mathcal{N})]_\theta = \tilde{I}_{s,p}^\eta(L_s^r MO(\mathcal{N})).$$

For $2 < p \leq \infty$, the following martingale norms were defined in [18]:

$$\|x\|_{L_p MO(\mathcal{N})} = \max\{\|x\|_{L_p^c MO(\mathcal{N})}, \|x\|_{L_p^r MO(\mathcal{N})}\},$$

respectively,

$$\|x\|_{H_p(\mathcal{N})} = \max\{\|x\|_{H_p^c(\mathcal{N})}, \|x^*\|_{H_p^c(\mathcal{N})}\}.$$

We now state the first version of our main result.

Theorem 3.17. *Let $2 < p < s < q \leq \infty$, $0 < \theta < 1$ such that $\frac{1}{s} = \frac{1-\theta}{q} + \frac{\theta}{p}$, and $0 \leq \eta \leq 1$. Then, with equivalent norms,*

$$[I_{q,p}^\eta(L_q MO(\mathcal{N})), L_p(\mathcal{N})]_\theta = I_{s,p}^\eta(L_s(\mathcal{N})).$$

Proof. Recall that, by Corollary 3.12, we have

$$\|x\|_{L_p^c MO(\mathcal{N})} \leq \lambda_p \|x\|_p,$$

where $\lambda_p = O(1)$ as $p \rightarrow \infty$. Combining this inequality with Corollary 3.15, Proposition 3.16 and Corollary 5.6 of [35], it follows that

$$\begin{aligned} [I_{q,p}^\eta(L_q MO(\mathcal{N})), L_p(\mathcal{N})]_\theta &\subseteq [\tilde{I}_{q,p}^\eta(L_q^c MO), L_p^c MO]_\theta \cap [\tilde{I}_{q,p}^\eta(L_q^r MO), L_p^r MO]_\theta \\ &\subseteq \tilde{I}_{s,p}^\eta(L_s^c MO) \cap \tilde{I}_{s,p}^\eta(L_s^r MO) \\ &= \tilde{I}_{s,p}^\eta(L_s^c MO \cap L_s^r MO) \\ &\subseteq \tilde{I}_{s,p}^\eta(L_s(\mathcal{N})), \end{aligned}$$

where the norm of the last inclusion above is $\lambda'_s \leq Cs$ as $s \rightarrow \infty$; here C is an absolute constant. The reverse inclusion follows by applying Kosaki's interpolation result (3.18), together with Corollary 3.15 and Proposition 3.16, namely

$$\begin{aligned} I_{s,p}^\eta(L_s(\mathcal{N})) &\subseteq [I_{q,p}^\eta(L_q(\mathcal{N})), L_p(\mathcal{N})]_\theta \\ &\subseteq [I_{q,p}^\eta(L_q^c MO \cap L_q^r MO), L_p(\mathcal{N})]_\theta \\ &= [I_{q,p}^\eta(L_q MO), L_p(\mathcal{N})]_\theta. \end{aligned} \quad \square$$

As an application, we obtain our main result of this section. The following continuous inclusion is proved using the fact that we have an interpolation scale of spaces.

Theorem 3.18. *Let $1 \leq p < \infty$ and $0 \leq \eta \leq 1$, then the inclusion map*

$$I_{\infty,p}^\eta(BMO) \subset L_p(\mathcal{N})$$

is bounded, with norm $c(p) \leq cp$, where c is an absolute constant.

Proof. Assume that $p \geq 8$. Let $4 < s < p/2 < \infty$ and define $0 < \theta < 1$ such that $\frac{1}{p} = \frac{1-\theta}{\infty} + \frac{\theta}{s}$. Since the complex interpolation method is an exact interpolation functor of exponent θ , we deduce from (the proof of) Theorem 3.17, that for a finite martingale difference sequence $x = (d_k)_{k=0}^n \in BMO$ we have

$$\begin{aligned} \|D^{\frac{1-\eta}{p}} x D^{\frac{\eta}{p}}\|_p &\leq c(p) \|D^{\frac{1-\eta}{s}} x D^{\frac{\eta}{s}}\|_{[I_{\infty,s}^\eta(BMO), L_s(\mathcal{N})]_\theta} \\ &\leq c(p) \|x\|_{BMO}^{1-\theta} \|D^{\frac{1-\eta}{s}} x D^{\frac{\eta}{s}}\|_s^\theta \leq c(p) \|x\|_{BMO}^{1-\theta} \|D^{\frac{1-\eta}{p}} x D^{\frac{\eta}{p}}\|_p^\theta, \end{aligned}$$

where $c(p) \leq Cp$, with C an absolute constant. Thus, we get

$$(3.44) \quad \|D^{\frac{1-\eta}{p}} x D^{\frac{\eta}{p}}\|_p \leq (c(p))^{\frac{1}{1-\theta}} \|x\|_{BMO}.$$

Under the hypotheses on p and s , a simple computation shows that

$$(c(p))^{\frac{1}{1-\theta}} \leq cp,$$

where c is an absolute constant. Therefore (3.44) implies that

$$\|D^{\frac{1-\eta}{p}} x D^{\frac{\eta}{p}}\|_p \leq cp \|x\|_{BMO}.$$

For an arbitrary element $x = (d_k)_{k \geq 0}$ in BMO , we consider the sequence of projections $P_n(x)$, where

$$P_n(x) = (d_k)_{k=0}^n,$$

whose images $I_{\infty,p}^\eta(P_n(x))$ are uniformly bounded and thus converge in $L_p(\mathcal{N})$ to $I_{\infty,p}^\eta(x)$. This yields the assertion for $p \geq 8$. Since $L_8(\mathcal{N})$ embeds into $L_p(\mathcal{N})$ for all $1 \leq p < 8$, the proof is complete. $\square$

Note that the symmetric embedding

$$\Psi_p(x) = D^{\frac{1}{2p}} x D^{\frac{1}{2p}}$$

is positivity preserving and thus, it can be considered as the natural embedding of $\mathcal{N}$ into $L_p(\mathcal{N})$. As a special case of Theorem 3.18 we obtain the following

Corollary 3.19. *If $1 \leq p < \infty$, then the inclusion map*

$$\Psi_p(BMO) \subset L_p(\mathcal{N})$$

is bounded, with norm $c(p) \leq cp$, where c is an absolute constant.

Remark 3.20. The following example which answers a question raised by Tao Mei, shows that the above inclusion does not hold if we only restrict to column, respectively row versions of the BMO norm. Let n be a positive integer and consider the von Neumann algebra

$$\mathcal{N} = L_\infty([0, 1]) \bar{\otimes} M_n,$$

where M_n is the algebra of $n \times n$ complex matrices. For $k \geq 1$ let Σ_k be the σ -algebra generated by dyadic intervals in $[0, 1]$ of length 2^{-k} . Denote by $\mathcal{N}_k$ the subalgebra $L_\infty([0, 1], \Sigma_k) \bar{\otimes} M_n$ of $\mathcal{N}$ and let $\mathcal{E}_k = \mathbb{E}_k \otimes \text{Id}_{M_n}$ be the conditional expectation onto $\mathcal{N}_k$. Let $\varepsilon_k \in \{-1, 1\}$ and define

$$x = \sum_{k=1}^n \varepsilon_k \otimes e_{1k}.$$

Then x is a martingale relative to the filtration $(\mathcal{N}_k)$. The martingale differences are given by $d_k(x) = \varepsilon_k \otimes e_{1k}$. A simple computation shows that $\|x\|_{L_p(\mathcal{N})} = n^{\frac{1}{2} - \frac{1}{p}}$, while

$$\|x\|_{BMO_c} = \sup_m \left\| \sum_{k=m}^n \mathcal{E}_m(d_k^*(x) d_k(x)) \right\|_\infty^{\frac{1}{2}} = \sup_k \|\varepsilon_k^2\|_\infty^{\frac{1}{2}} = 1.$$

Assume that $p > 2$. Then for any $C > 0$, there exists $n \geq 1$ such that $n^{\frac{1}{2}-\frac{1}{p}} > C$. This implies that $BMO_c(\mathcal{N})$ is not contained in $L_p(\mathcal{N})$.

4. MAIN RESULTS

In this section, we will show how to derive the John-Nirenberg type result from Theorem 3.18. Let us start with an immediate application.

Corollary 4.1. *Let $1 \leq p < \infty$, a positive integer n and an element $a \in L_p(\mathcal{N}_n)$. Then, there exists an absolute constant $c > 0$ such that for all $x \in \mathcal{N}$,*

$$(4.45) \quad \|(x - x_{n-1})a\|_p \leq cp \|x\|_{BMO} \|a\|_p.$$

Proof. Assume that $p \geq 2$. It suffices to show (4.45) for positive elements $a \in L_p(\mathcal{N}_n)$ with $\|a\|_p = 1$. Furthermore, by approximation with elements of the form $(a^p + \varepsilon D)^{\frac{1}{p}}$, we may assume that a has full support. Define a new state ϕ_a on $\mathcal{N}$ by

$$\phi_a(x) = \text{tr}(a^p x), \quad x \in \mathcal{N}.$$

Denote by $L_p(\mathcal{N}, \phi_a)$ the noncommutative L_p space associated to the state ϕ_a . Note that

$$(4.46) \quad \|x - x_{n-1}\|_{L_p(\mathcal{N}, \phi_a)} = \|(x - x_{n-1})a\|_p.$$

For all non-negative integers k define $\tilde{\mathcal{N}}_k = \mathcal{N}_{n+k}$. Then $\tilde{\mathcal{N}}_0 = \mathcal{N}_n$ and $(\tilde{\mathcal{N}}_k)_{k \geq 0}$ is a filtration of $\mathcal{N}$. Moreover, if $m \geq n$ and $x \in \mathcal{N}$, then $xa^p \in L_1(\mathcal{N}_m)$. Hence,

$$(4.47) \quad \phi_a(yx) = \text{tr}(a^p yx) = \text{tr}(yxa^p) = \text{tr}(\mathcal{E}_m(y)xa^p) = \phi_a(\mathcal{E}_m(y)x),$$

for all $y \in \mathcal{N}_m$. Denote by $BMO((\tilde{\mathcal{N}}_k)_{k \geq n}, \phi_a)$ the BMO space associated to the filtration $(\tilde{\mathcal{N}}_k)_{k \geq 0}$ of $\mathcal{N}$ and the state ϕ_a . From Theorem 3.18 it follows that

$$(4.48) \quad \|x - x_{n-1}\|_{L_p(\mathcal{N}, \phi_a)} \leq cp \|x - x_{n-1}\|_{BMO((\tilde{\mathcal{N}}_k)_{k \geq n}, \phi_a)},$$

The modular automorphism group of ϕ_a is given by

$$\sigma_t^{\phi_a}(x) = a^{ipt} x a^{-ipt}, \quad t \in \mathbb{R}.$$

From (4.47) we deduce that for all $k \geq 0$, the conditional expectations $\tilde{\mathcal{E}}_k = \mathcal{E}(\cdot | \tilde{\mathcal{N}}_k)$ associated with the state ϕ_a are given by the original conditional expectations, namely, $\tilde{\mathcal{E}}_k = \mathcal{E}_{k+n}$. Therefore,

$$\tilde{\mathcal{E}}_k(x - x_{n-1}) = \mathcal{E}_{k+n}(x - x_{n-1}) = x_{k+n} - x_{n-1}.$$

Hence, denoting $x - x_{n-1}$ by y , we obtain

$$\tilde{\mathcal{E}}_k((y - \tilde{\mathcal{E}}_{k-1}(y))^*(y - \tilde{\mathcal{E}}_{k-1}(y))) = \mathcal{E}_{n+k}((x - x_{n+k-1})^*(x - x_{n+k-1})).$$

By the definition of the BMO norm, it follows that

$$\|x - x_{n-1}\|_{BMO((\tilde{\mathcal{N}}_k)_{k \geq n}, \phi_a)} \leq \|x\|_{BMO}.$$

Together with (4.48) and (4.46), this yields the assertion in the case $p \geq 2$. Furthermore, if $1 \leq p < 2$ and $0 \leq a \in L_p(\mathcal{N})$ with $\|a\|_p = 1$, consider $a_1 = a^{\frac{1}{2}}$. Then $0 \leq a_1 \in L_{2p}(\mathcal{N})$ and $\|a_1\|_{2p} = 1$. By Hölder's inequality it follows that

$$(4.49) \quad \|(x - x_{n-1})a\|_p \leq \|(x - x_{n-1})a_1\|_{2p}.$$

Note that $2p \geq 2$. Applying (4.45) and (4.49) to a_1 , we obtain the conclusion. $\square$

Remark 4.2. Note that for any integer $n \geq 0$, the inclusion $\mathcal{N} \subset L_\infty^c(\mathcal{N}, \mathcal{E}_n)$ is injective. Thus, given $2 \leq p < \infty$, we can consider the space $[\mathcal{N}, L_\infty^c(\mathcal{N}, \mathcal{E}_n)]^{\frac{2}{p}}$ obtained by the upper method of complex interpolation of exponent $\frac{2}{p}$. These are particular examples of conditional L_p -spaces considered in [16]. Following ideas of Pisier [30], it is proved in [16] that

$$(4.50) \quad \|x\|_{[\mathcal{N}, L_\infty^c(\mathcal{N}, \mathcal{E}_n)]^{\frac{2}{p}}} = \sup_{\|a\|_{L_p(\mathcal{N}_n)} \leq 1} \|xa\|_{L_p(\mathcal{N})}.$$

Therefore, as already mentioned in the introduction, the BMO_p^c norms defined by (1.9) are in fact interpolation norms, namely

$$(4.51) \quad \|x\|_{BMO_p^c} = \|x - x_{n-1}\|_{[\mathcal{N}, L_\infty^c(\mathcal{N}, \mathcal{E}_n)]^{\frac{2}{p}}}.$$

Proof of Theorem 1.1. Let $2 < p < \infty$. Trivially, we have

$$[\mathcal{N}, L_\infty^c(\mathcal{N}, \mathcal{E}_n)]^{\frac{2}{p}} \subset L_\infty^c(\mathcal{N}, \mathcal{E}_n).$$

Therefore, we deduce that

$$\|\mathcal{E}_n((x - x_{n-1})^*(x - x_{n-1}))\|_\infty^{\frac{1}{2}} \leq \|x - x_{n-1}\|_{[\mathcal{N}, L_\infty^c(\mathcal{N}, \mathcal{E}_n)]^{\frac{2}{p}}}.$$

Combining this with (4.51), it follows immediately that

$$\|x\|_{BMO} \leq \|x\|_{BMO_p}.$$

For the converse, we deduce from (1.9) and Corollary 4.1 that

$$\|x\|_{BMO_p^c} = \sup_{\|a\|_{L_p(\mathcal{N}_n)} \leq 1} \|(x - x_{n-1})a\|_p \leq cp \|x\|_{BMO}.$$

Applying the same argument to x^* , we deduce that

$$\|x\|_{BMO_p} \leq cp \|x\|_{BMO},$$

which concludes the proof. $\square$

Remark 4.3. As an application of Theorem 1.1 we now prove that the noncommutative analogue of the classical result

$$(4.52) \quad BMO \subset L_{\exp}$$

holds in the setting of a semifinite von Neumann algebra $\mathcal{N}$ equipped with a normal, faithful trace τ . In this context the space $L_{\exp}(\mathcal{N})$ can be defined following the general scheme of *symmetric spaces* associated to $(\mathcal{N}, \tau)$ and a rearrangement invariant Banach function space developed in [6] and [7]. Therefore, motivated by the classical definition of the Zygmund space $L_{\exp}$, we define

$$(4.53) \quad \|x\|_{L_{\exp}(\mathcal{N})} = \inf\{\lambda > 0 : \tau(e^{\frac{|x|}{\lambda}-1}) \leq 1\}.$$

Suppose that $\|x\|_{BMO} \leq 1$ and let $\lambda > 0$. Then, using the power series expansion for the exponential, it follows from Theorem 1.1 that

$$\begin{aligned} \tau(e^{\frac{|x|}{\lambda}-1}) &= \frac{1}{e} \sum_{k=0}^{\infty} \frac{\tau(|x|^k)}{\lambda^k k!} = \frac{1}{e} \left(1 + \sum_{k=1}^{\infty} \frac{\|x\|_k^k}{\lambda^k k!} \right) \\ &\leq \frac{1}{e} \left(1 + \sum_{k=1}^{\infty} \frac{c^k k^k}{\lambda^k k!} \right) \\ &\leq \frac{1}{e} \left(1 + \sum_{k=1}^{\infty} \frac{c^k e^k}{\lambda^k} \right) = \frac{1}{e} \sum_{k=0}^{\infty} \left(\frac{ce}{\lambda} \right)^k. \end{aligned}$$

For the last inequality we have used the fact, which can be verified by induction, that $k^k \leq k!e^k$, for all integers $k \geq 1$. A simple computation shows that if $\frac{1}{\lambda} \leq \frac{1}{ce} (1 - \frac{1}{e})$, then $\frac{1}{e} \sum_{k=0}^{\infty} \left(\frac{ce}{\lambda} \right)^k \leq 1$.

By (4.53), it follows that

$$\|x\|_{L_{\exp}(\mathcal{N}, \tau)} \leq \frac{1}{ce} \left(1 - \frac{1}{e} \right).$$

Hence $x \in L_{\exp}(\mathcal{N})$ and (4.52) is proved.

Open problem: Is there a universal constant C such that if $x \in BMO$, then

$$\sup_n \|\mathcal{E}_n(|x - x_{n-1}|^p)\|_{\infty}^{\frac{1}{p}} \leq Cp \|x\|_{BMO},$$

for all $1 \leq p < \infty$?

The noncommutative versions of function space BMO have been developed by Mei [25]. Namely, for a function $x : \mathbb{R} \rightarrow \mathcal{N}$, define the following norm

$$\|x\|_{BMO_c} = \sup_{\emptyset \neq I} \left\| \int_I (x(t) - x_I)^* (x(t) - x_I) \frac{dt}{|I|} \right\|_{\mathcal{N}},$$

where $x_I = \int_I x(t) \frac{dt}{|I|}$, for every nonempty interval I . Furthermore, define

$$\|x\|_{BMO_r} = \|x^*\|_{BMO_c}$$

and

$$\|x\|_{BMO} = \max\{\|x\|_{BMO_c}, \|x\|_{BMO_r}\}.$$

As in the martingale setting, we use interpolation to define the BMO_p -norms in this context. Namely, for $2 < p < \infty$ and a fixed nonempty interval I , let $\mu_I = \frac{dt}{|I|}$ and define the space

$$\mathcal{N}(L_p^c(\mu_I)) = [\mathcal{N} \bar{\otimes} L_\infty(\mu_I), \mathcal{N} \bar{\otimes} L_2^c(\mu_I)]^{\frac{2}{p}}$$

by (the upper method of) complex interpolation. Recall that $\mathcal{N} \bar{\otimes} L_\infty(\mu_I) = L_\infty(I, \mu_I; \mathcal{N})$ is the usual space of measurable, essentially bounded $\mathcal{N}$ -valued functions, while the space $\mathcal{N} \bar{\otimes} L_2^c(\mu_I)$ is defined following the operator space tradition of viewing $L_2(\mu_I)$ implemented as a column in $B(L_2(\mu_I))$. Equivalently,

$$\|x\|_{\mathcal{N} \bar{\otimes} L_2^c(\mu_I)} = \left\| \int_I (x(t))^* x(t) d\mu_I(t) \right\|_{\mathcal{N}}^{\frac{1}{2}}$$

for all $\mathcal{N}$ -valued functions which are measurable with respect to the strong operator topology. Note that $\mathcal{E}_I(x) = \int_I x(t) d\mu_I(t)$ is a conditional expectation onto $\mathcal{N}_I = \mathcal{N} \bar{\otimes} L_\infty(\mu_I)$. Therefore, the spaces $\mathcal{N}(L_p^c(\mu_I))$ are special cases of conditional L_p -spaces considered in [16]. In particular the following norm estimate holds

$$(4.54) \quad \|1_I x\|_{\mathcal{N}(L_p^c(\mu_I))} = \sup_{\|a\|_{L_p(\mathcal{N})} \leq 1} \left(\int_I \|x(t)a\|_{L_p(\mathcal{N})}^p d\mu_I(t) \right)^{\frac{1}{p}}.$$

As in [25], we will use the following result to transfer martingale results to intervals.

Lemma 4.4 (T. Mei). *Let $I \subset \mathbb{R}$ be an interval. Then there exists a dyadic interval $J = [\frac{k-1}{2^n}, \frac{k}{2^n}]$ or $J' = [\frac{1}{3 \cdot 2^{n'}} + \frac{k'-1}{2^{n'}}, \frac{1}{3 \cdot 2^{n'}} + \frac{k'}{2^{n'}}]$ such that*

$$I \subset J \quad \text{and} \quad |J| \leq 6|I|$$

or

$$I \subset J' \quad \text{and} \quad |J'| \leq 6|I|.$$

Lemma 4.5. *For a nonempty interval $I \subset \mathbb{R}$ and an element $x \in \mathcal{N} \bar{\otimes} L_\infty(\mathbb{R})$ let*

$$\|x\|_{p,I} = \sup_{\|a\|_{L_p(\mathcal{N})} \leq 1} \|(1_I x - x_I)a\|_{L_p(I, \mu_I; L_p(\mathcal{N}))}.$$

Then, for all intervals $I \subset J \subset \mathbb{R}$,

$$\|x\|_{p,I} \leq 2 \left(\frac{|J|}{|I|} \right)^{\frac{1}{p}} \|x\|_{p,J}.$$

Proof. This is of course a standard argument. Let $a \in L_p(\mathcal{N})$. Then we have

$$\|(x_I - x_J)a\|_p = \left\| \int_I [(x(s) - x_J)a] d\mu_I(s) \right\|_p \leq \left(\int_I \|(x(t) - x_J)a\|_p^p d\mu_I(t) \right)^{\frac{1}{p}}.$$

Therefore, we obtain

$$\begin{aligned}
\|(1_I x - x_I)a\|_{L_p(I, \mu_I; L_p(\mathcal{N}))} &\leq \|(1_I x - x_J)a\|_{L_p(I, \mu_I; L_p(\mathcal{N}))} + \|(x_I - x_J)a\|_p \\
&\leq 2 \left(\int_I \|(x(t) - x_J)a\|_p^p d\mu_I(t) \right)^{\frac{1}{p}} \\
&\leq 2 \left(\frac{|J|}{|I|} \right)^{\frac{1}{p}} \left(\int_J \|(x(t) - x_J)a\|_p^p d\mu_J(t) \right)^{\frac{1}{p}}.
\end{aligned}$$

Taking supremum over all elements a in the unit ball of $L_p(\mathcal{N})$ implies the assertion. $\square$

We are now ready to formulate a noncommutative interval version of the John-Nirenberg theorem.

Theorem 4.6. *For all $2 < p < \infty$, there exists an absolute constant $c > 0$ such that*

$$\|x\|_{BMO} \leq \sup_{\emptyset \neq I \text{ interval}} \max\{\|x - x_I\|_{\mathcal{N}(L_p^c(\mu_I))}, \|x^* - x_I^*\|_{\mathcal{N}(L_p^c(\mu_I))}\} \leq cp \|x\|_{BMO}.$$

Proof. Denote by $\tilde{\mathcal{N}}$ the von Neumann algebra tensor product $\mathcal{N} \bar{\otimes} L_\infty[0, 1]$. For each integer $n \geq 0$, let Σ_n be the σ -algebra generated by dyadic intervals in $[0, 1]$ of length 2^{-n} . Denote by $\tilde{\mathcal{N}}_n$ the subalgebra $\mathcal{N} \bar{\otimes} L_\infty([0, 1], \Sigma_n)$ of $\tilde{\mathcal{N}}$ and let $\tilde{\mathcal{E}}_n$ be the conditional expectation onto $\tilde{\mathcal{N}}_n$. Consider the BMO space associated to the filtration $(\tilde{\mathcal{N}}_n)_{n \geq 0}$ of $\tilde{\mathcal{N}}$. For $x \cong (x_n)_{n \geq 0}$, where $x_n = \tilde{\mathcal{E}}_n(x)$, we deduce that

$$\sup_n \max\{\|(x - x_n)\|_{L_p^c(\tilde{\mathcal{N}}, \tilde{\mathcal{E}}_n)}, \|(x - x_n)^*\|_{L_p^c(\tilde{\mathcal{N}}, \tilde{\mathcal{E}}_n)}\} \leq \|x\|_{BMO}.$$

An application of (4.54) shows that

$$\begin{aligned}
\|x - x_n\|_{L_p^c(\tilde{\mathcal{N}}, \tilde{\mathcal{E}}_n)}^p &= \sup_{\sum_k \|a_k\|_p^p |I_k| \leq 1} \sum_k |I_k| \int_{I_k} \|(x(t) - x_{I_k})a_k\|_p^p d\mu_{I_k}(t) \\
&= \sup_k \|1_{I_k} x - x_{I_k}\|_{\mathcal{N}(L_p^c(\mu_{I_k}))}^p.
\end{aligned}$$

Here the supremum is taken over all intervals $I_k \subset [0, 1]$ of length 2^{-n} . Therefore, we may reformulate the martingale result in Theorem 1.1 as

$$\begin{aligned}
&\sup_{I \text{ dyadic } \subset [0, 1]} \max\{\|1_I x - x_I\|_{\mathcal{N}(L_p^c(\mu_I))}, \|(1_I x - x_I)^*\|_{\mathcal{N}(L_p^c(\mu_I))}\} \\
&\leq cp \sup_{I \text{ dyadic } \subset [0, 1]} \max\{\|1_I x - x_I\|_{\mathcal{N}(L_2^c(\mu_I))}, \|(1_I x - x_I)^*\|_{\mathcal{N}(L_2^c(\mu_I))}\}.
\end{aligned}$$

Using the dilation and translation invariance properties of the Lebesgue measure, it follows for all integers $k \geq 0$ that

$$\begin{aligned} & \sup_{I \text{ dyadic } \subset [2^{-k}, 2^k]} \max\{\|1_I x - x_I\|_{\mathcal{N}(L_p^c(\mu_I))}, \|(1_I x - x_I)^*\|_{\mathcal{N}(L_p^c(\mu_I))}\} \\ & \leq cp \sup_{I \text{ dyadic } \subset [2^{-k}, 2^k]} \max\{\|1_I x - x_I\|_{\mathcal{N}(L_2^c(\mu_I))}, \|(1_I x - x_I)^*\|_{\mathcal{N}(L_2^c(\mu_I))}\}. \end{aligned}$$

Taking the supremum over k , we obtain the desired inequality for arbitrary dyadic martingales. Applying Mei's result 4.4, together with Lemma 4.5 the assertion follows. $\square$

We now discuss an analogue of the classical large deviation inequality (1.12) in this setting. Our main tool is the following corollary to a noncommutative version of Chebychev inequality proved by Defant and Junge (see [8]).

Corollary 4.7 (Defant-Junge). *Let $2 \leq p < \infty$ and $x \in L_p(\mathcal{N})$. Then, for every $\varepsilon > 0$, there exists a projection $f \in \mathcal{N}$ with $\phi(1 - f) \leq \varepsilon$, such that whenever $yD^{\frac{1}{p}} = wx$, where $w, y \in \mathcal{N}$, then*

$$(4.55) \quad \sup_n \|f \mathcal{E}_n(y^* y) f\|^{\frac{1}{2}} \leq (1 + \sqrt{10}) \varepsilon^{-\frac{1}{p}} \|w\|_{\infty} \|x\|_p.$$

Moreover,

$$(4.56) \quad \sup_n \|\mathcal{E}_n(y) f\| \leq (1 + \sqrt{10}) \varepsilon^{-\frac{1}{p}} \|w\|_{\infty} \|x\|_p.$$

Proof of Theorem 1.2. Recall that the state ϕ satisfies condition (1.8). Let $c_2 = e$. Furthermore, consider $t \geq \frac{1}{c_1}$ and let $\varepsilon := e^{-tc_1}$, where c_1 is a constant to be made precise later. Define $p = 4 \ln(\varepsilon^{-1}) = 4tc_2$. By construction it follows that $4 \leq p < \infty$. Now let $x \in BMO$ with $\|x\|_{BMO} \leq 1$. Denote by y the element $(x - x_0)D^{\frac{1}{p}}$. Note that $y \in L_p(\mathcal{N})$. By the noncommutative Doob's inequality (see [14]) and Theorem 1.1, we get

$$(4.57) \quad \left\| \sup_n \mathcal{E}_n(y^* y) \right\|_{\frac{p}{2}} \leq d_{\frac{p}{2}} \|y\|_p^2 \leq d_{\frac{p}{2}}^2 c^2 p^2 \|x - x_0\|_{BMO}^2,$$

where c is the universal constant in Theorem 1.1 and $d_{\frac{p}{2}}$ is the constant in the noncommutative Doob's inequality. Since $\frac{p}{2} \geq 2$, it follows that $d_{\frac{p}{2}} = 2$ (see [14]). Furthermore, note that

$$(4.58) \quad \|x - x_0\|_{BMO} \leq \|x\|_{BMO}.$$

Indeed, we have the estimates

$$\mathcal{E}_0((x - x_0)^*(x - x_0)) = \mathcal{E}_0(x^* x - x^* x_0 - \mathcal{E}_0(x^*)x + \mathcal{E}_0(x^*)\mathcal{E}_0(x)) = \mathcal{E}_0(x^* x) - \mathcal{E}_0(x^*)\mathcal{E}_0(x).$$

By Kadison's inequality it follows that

$$(4.59) \quad \|\mathcal{E}_0((x - x_0)^*(x - x_0))\| \leq \|\mathcal{E}_0(x^* x)\|.$$

Moreover, for $n \geq 1$ we have

$$(x - \mathcal{E}_0(x)) - \mathcal{E}_{n-1}(x - \mathcal{E}_0(x)) = (x - \mathcal{E}_0(x)) - (\mathcal{E}_{n-1}(x) - \mathcal{E}_0(x)) = x - \mathcal{E}_{n-1}(x).$$

Combining this with (4.59) we deduce that

$$\|x - x_0\|_{BMO_c} \leq \|x\|_{BMO_c},$$

which yields the inequality (4.58).

A further application of Kadison's inequality and (4.57), together with (4.58) implies that

$$\left\| \sup_m \mathcal{E}_m(y^*) \mathcal{E}_m(y) \right\|_{\frac{p}{2}} \leq 2c^2 p^2.$$

Then by [14], Remark 3.7, there exists $0 \leq a \in L_p(\mathcal{N})$ and a sequence of positive contractions $w_m \in \mathcal{N}$ such that

$$\mathcal{E}_m(y^*) \mathcal{E}_m(y) = a w_m a, \quad \|a\|_p^2 \sup_m \|w_m\|_\infty = \left\| \sup_m \mathcal{E}_m(y^*) \mathcal{E}_m(y) \right\|_{\frac{p}{2}} \leq \sqrt{2} c p.$$

Using the polar decomposition of $\mathcal{E}_m(y)$, we find partial isometries u_m such that, denoting $u_m w_m^{\frac{1}{2}}$ by z_m , we obtain

$$(4.60) \quad \mathcal{E}_m(y) = z_m a, \quad \|a\|_p \sup_m \|z_m\|_\infty \leq \sqrt{K} p.$$

Note that $\mathcal{E}_m(x) - x_0 \in \mathcal{N}$, for all non-negative integers m . Indeed, for all $n \geq 0$,

$$1 \geq \|x\|_{BMO_c}^2 = \sup_n \left\| \sum_{k \geq n} \mathcal{E}_n(d_k^* d_k) \right\|_\infty \geq \|\mathcal{E}_n(d_n^* d_n)\|_\infty = \|d_n^* d_n\|_\infty.$$

This implies that $d_n^* d_n \in \mathcal{N}$ and hence $d_n \in \mathcal{N}$. Therefore $\mathcal{E}_m(x) - x_0 = \sum_{k=1}^m d_k \in \mathcal{N}$. Since

$$\mathcal{E}_m(y) = (\mathcal{E}_m(x) - x_0) D^{\frac{1}{p}},$$

we are now in the condition of applying Corollary 4.7 and deduce the existence of a projection $f \in \mathcal{N}$ with $\phi(1 - f) \leq \varepsilon$ such that for all $m \geq 0$,

$$\|(\mathcal{E}_m(x) - x_0) f\| \leq (1 + \sqrt{10}) \varepsilon^{-\frac{1}{p}} \|z_m\|_\infty \|a\|_p.$$

Using the estimates (4.60) we deduce that

$$\|(\mathcal{E}_m(x) - x_0) f\| \leq K p \varepsilon^{-\frac{1}{p}},$$

where $K = (1 + \sqrt{10}) \sqrt{2} c$. Therefore $\|(\mathcal{E}_m(x) - x_0) h\| \leq K p \varepsilon^{-\frac{1}{p}}$, for all $h \in fH$. Since the sequence $(\mathcal{E}_m)_{m \geq 0}$ converges to $\text{Id}_{\mathcal{N}}$ in the strong operator topology, it follows that

$$(4.61) \quad \|(x - x_0) f\| \leq K p \varepsilon^{-\frac{1}{p}}.$$

We now make precise the constant c_1 . Namely, let

$$c_1 := \frac{1}{4K e^{1/4}} = \frac{1}{4\sqrt{2}(1 + \sqrt{10}) c e^{1/4}}.$$

A simple computation yields the equality $K p \varepsilon^{-\frac{1}{p}} = t$. By (4.61) we deduce the inequality $\|(x - x_0) f\| \leq t$. Moreover, $\phi(1 - f) \leq \varepsilon < c_2 e^{-t c_1}$, which yields the assertion.

For $0 < t < \frac{1}{c_1}$, let $f = 0$. The condition $\|(x - x_0)f\| \leq t$ is then automatically satisfied. Furthermore, note that

$$\phi(1 - f) = \phi(1) = 1 < \frac{e}{e^{tc_1}} = c_2 e^{-tc_1}.$$

This completes the proof. $\square$

Remark 4.8. After completing this paper, we have been informed that Tao Mei has obtained a simple proof of the inequalities (1.11), but only in the tracial setting, based on the interpolation result proved in [26].

REFERENCES

- [1] R.F. BASS: *Probabilistic techniques in analysis*, Springer-Verlag, New York 1995.
- [2] D. BURKHOLDER: *Martingales and singular integrals in Banach spaces*, Handbook on the Geometry of Banach Spaces, Volume 1, W.B. Johnson and J. Lindenstrauss editors, North Holland (2001), 233-269.
- [3] J. BERGH, J. LÖFSTRÖM: *Interpolation Spaces. An Introduction*, Springer-Verlag, New-York 1976.
- [4] E. A. CARLEN AND P. KRÉE, *On martingale inequalities in noncommutative stochastic analysis*, J. Funct. Analysis **158** (1998), 475-508.
- [5] C. BENNETT, R. SHARPLEY: *Interpolation of operators*, Academic Press 1988.
- [6] P.G. DODDS, T.K. DODDS, B. DE PAGTER: *Non-commutative Banach function spaces*, Math. Z. **201** (1989), 583-597.
- [7] P.G. DODDS, T.K. DODDS, B. DE PAGTER: *Fully symmetric operator spaces*, Integr. Eq. Op. Theory Vol **15** (1992), 942-972.
- [8] A. DEFANT, M. JUNGE: *Riesz summation in noncommutative L_p -spaces*, Preprint.
- [9] J. GARCIA-CUERVA, J. RUBIO DE FRANCIA: *Weighted norm inequalities and related topics*, North Holland Mathematics Studies **116**, Notas de Matematica 104, Elsevier Sci. Publishers B.V., 1985.
- [10] J.B. GARNETT: *Bounded analytic functions*, Academic Press, Inc. (London) Ltd., 1981.
- [11] A. GARSIA: *Martingale Inequalities: Seminar Notes on Recent Progress*, Math. Lecture Notes Series, 1973.
- [12] N. GHOUSSEUB, G. GODEFROY, B. MAUREY, W. SCHACHERMAYER: *Some topological and geometric structures in Banach spaces*, Memoirs Amer. Math. Soc. **379** (1987).
- [13] F. JOHN, L. NIRENBERG: *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. **167** (1961), 416-426.
- [14] M. JUNGE, *Doob's inequality for non-commutative martingales*, J. reine angew. Math. **549** (2002), 149-190.
- [15] M. JUNGE, *Fubini's theorem for ultraproducts of noncommutative L_p spaces*; to appear in Canadian J. Math.
- [16] M. JUNGE, J. PARCET, *Interpolation of amalgamated non-commutative L_p spaces*, Preprint.
- [17] M. JUNGE, D. SHERMAN, *Noncommutative L_p modules*, To appear in J. Operator Theory.
- [18] M. JUNGE AND Q. XU, *Non-commutative Burkholder/Rosenthal inequalities*, Ann. Probab. **31** (2003) no. 2, 948-995.
- [19] M. JUNGE AND Q. XU, *Non-commutative Burkholder/Rosenthal inequalities: Applications*, In preparation.
- [20] M. JUNGE, Q. XU: *The optimal orders of growth of the best constants in some non-commutative martingale inequalities*, Preprint, 2001.
- [21] R.V. KADISON, J.R. RINGROSE: *Fundamentals of the Theory of Operator Algebras I, II*, Academic Press, 1986.
- [22] P. KOOSIS: *INTRODUCTION TO H_p -SPACES*, Cambridge tracts in mathematics **115**, Cambridge University Press 1998.

- [23] H. KOSAKI: *Applications of the complex interpolation method to a von Neumann algebra: Non-commutative L_p -spaces*, J. Funct. Analysis **56** (1984), 29-78.
- [24] E.C. LANCE: *Hilbert C^* -modules. A toolkit for operator algebraists*, London Math. Soc. Lecture Notes Series 210, Cambridge Univ. Press, Cambridge, 1995.
- [25] T. MEI: *Operator-valued Hardy spaces*, Preprint.
- [26] M. MUSAT: *Interpolation between non-commutative BMO and non-commutative L_p -spaces*, J. Funct. Analysis **202** (2003), 195-225.
- [27] E. NELSON: *Notes on non-commutative integration*, J. Funct. Analysis **15** (1974), 103-116.
- [28] G. PEDERSEN, M. TAKESAKI: *The Radon-Nikodym theorem for von Neumann algebras*, Acta Math. **130** (1973), 53-87.
- [29] K.E. PETERSEN: *Brownian motion, Hardy spaces and bounded mean oscillation*, London Math. Soc. Lecture Notes Series 28, Cambridge Univ. Press, Cambridge 1977.
- [30] G. PISIER: *Projections from a von Neumann algebra onto a subalgebra*, Bull. Soc. Math. France **123** (1995), 139-153.
- [31] G. PISIER: *The operator Hilbert space OH , complex interpolation and tensor norms*, Mem. Amer. Math. Soc. **122** (1996), no. 595.
- [32] G. PISIER, Q. XU: *Non-commutative martingale inequalities*, Comm. Math. Phys. **189** (1997), no. 3, 667-698.
- [33] G. PISIER, Q. XU: *Non-commutative L_p -spaces, Handbook of the Geometry of Banach Spaces II*, Ed. W.B. Johnson, J. Lindenstrauss, North Holland (2003), 1459-1517.
- [34] N. RANDRIANANTOANINA: *Non-commutative martingale transforms*, J. Funct. Analysis **194** (2002), 181-212.
- [35] N. RANDRIANANTOANINA: *A weak type inequality for non-commutative martingales and applications*, Preprint.
- [36] M.A. RIEFFEL: *Induced representations of C^* -algebras*, Adv. Math. **13** (1974), 176-257.
- [37] E.M. STEIN: *Topics in harmonic analysis related to the Littlewood-Paley theory*, Princeton University Press, Princeton, NJ, 1970.
- [38] E.M. STEIN: *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, NJ, 1970.
- [39] S. STRATILA: *Modular Theory in Operator Algebras*, Abacus Press, Tunbridge Wells, Kent, England, 1981.
- [40] S. STRATILA, L. ZSIDO: *Lectures on von Neumann algebras*, Abacus Press, Tunbridge Wells, Kent, England, 1971.
- [41] M. TAKESAKI: *Theory of Operator Algebras I*, Springer-Verlag, New-York, 1979.
- [42] M. TAKESAKI: *Conditional expectations in von Neumann algebras*, J. Funct. Analysis **9** (1972), 306-321.
- [43] M. TERP: *L^p -spaces associated with von Neumann algebras I and II*, Copenhagen Univ. 1981.

⁽¹⁾DEPARTMENT OF MATHEMATICS, 1409 WEST GREEN STREET, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801.

E-mail address: `junge@math.uiuc.edu`

⁽²⁾DEPARTMENT OF MATHEMATICS, 0112, 9500 GILMAN DRIVE, UNIVERSITY OF CALIFORNIA , SAN DIEGO, LA JOLLA, CA 92093-0112.

E-mail address: `mmusat@math.ucsd.edu`